

FULL-LENGTH REVIEW – Food Microbiology

The forgotten role of food cultures

François Bourdichon^{1,2,*†}, Emmanuelle Arias³, Andrzej Babuchowski⁴, Anne Bückle⁵, Fabio Dal Bello⁶, Aurélie Dubois⁷, Alessandra Fontana², Duresa Fritz⁸, Rober Kemperman⁹, Svend Laulund¹⁰, Olivia McAuliffe¹¹, Marta Hanna Miks^{12,13}, Photis Papademas¹⁴, Vania Patrone², Dinesh K. Sharma¹⁵, Edward Sliwinski¹⁶, Catherine Stanton¹¹, Ueli Von Ah³, Su Yao¹⁷ and Lorenzo Morelli²

¹Food Safety, Microbiology, Hygiene, 16 Rue Gaston de Caillavet, 75015 Paris, France, ²Facoltà di Scienze agrarie, alimentari ambientali, Università Cattolica del Sacro Cuore, Via Emilia Parmense, Piacenza-Cremona, Italy, ³AGROSCOPE, Food Microbial Systems, Schwarzenburgstrasse 161, CH-3003 Bern, Switzerland, ⁴Dairy Industry Innovation Institute, Kormoranow St., Mragowo 11–700, Poland, ⁵Milchprüfing Baden-Württemberg e.V., Marie-Curie-Straße 19, 73230 Kirchheim, u.T., Germany, ⁶Sacco S.r.l., Via Alessandro Manzoni, Cadorago (Co), Italy, ⁷International Dairy Federation, 70 Boulevard Auguste Reyers, 1030 Brussels, Belgium, ⁸International Flavors and Fragrances, 20 rue Brunel, Paris 75017, France, ⁹Lesaffre International, 152 rue du Docteur Yersin, 59120 Loos, France, ¹⁰Chr. Hansen A/S, Agern Allé 24, 2970 Hoersholm, Denmark, ¹¹Teagasc Food Research Centre, Moorepark, Fermoy, Cork, Ireland, ¹²Glycom A/S, Kogle Allé 4, 2970 Hørsholm, Denmark, ¹³Faculty of Food Science, Food Biochemistry, University of Warmia and Mazury in Olsztyn, Plac Cieszyński 1, 10–726 Olsztyn, Poland, ¹⁴Department of Agricultural Sciences, Biotechnology and Food Science, Cyprus University of Technology, Archiepiskopou Kyprianou, PO BOX 50329, Limassol, Cyprus, ¹⁵National Dairy Development Board, Anand 388001, India, ¹⁶The European Federation of Food Science & Technology, Nieuwe Kanaal 9a, 6709 PA, Wageningen, The Netherlands and ¹⁷China National Research Institute of Food & Fermentation Industries, China Center of Industrial Culture Collection, Building 6, No.24, Jiuxianqiaozhong Road, Chaoyang District, Beijing 100015, PR China

*Corresponding author: Food Safety, Microbiology, Hygiene, 16 Rue Gaston de Caillavet, 75015 Paris, France. Tel: +33 6 24 75 21 49;

E-mail: francois.bourdichon@gmail.com

One sentence summary: Perspective on the potential of food cultures for the biopreservation of food products and applications to limit food waste.

Editor: Egon Hansen

[†]François Bourdichon, <http://orcid.org/0000-0003-0373-0407>

ABSTRACT

Fermentation is one of if not the oldest food processing technique, yet it is still an emerging field when it comes to its numerous mechanisms of action and potential applications. The effect of microbial activity on the taste, bioavailability and

Received: 14 August 2020; Accepted: 1 July 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

preservation of the nutrients and the different food matrices has been deciphered by the insights of molecular microbiology. Among those roles of fermentation in the food chain, biopreservation remains the one most debated. Presumably because it has been underestimated for quite a while, and only considered – based on a food safety and technological approach – from the toxicological and chemical perspective. Biopreservation is not considered as a traditional use, where it has been by design – but forgotten – as the initial goal of fermentation. The ‘modern’ use of biopreservation is also slightly different from the traditional use, due mainly to changes in cooling of food and other ways of preservation. Extending shelf life is considered to be one of the properties of food additives, classifying – from our perspective – biopreservation wrongly and forgetting the role of fermentation and food cultures. The present review will summarize the current approaches of fermentation as a way to preserve and protect the food, considering the different way in which food cultures and this application could help tackle food waste as an additional control measure to ensure the safety of the food.

Keywords: food cultures; fermentation; biopreservation; food safety; regulation; mechanism of action

INTRODUCTION

Food preservation has been a key concern since the earliest days of humanity. Among the numerous empirical processes that have been developed and passed down, fermentation is one of the oldest preservation techniques and still widely used in various food matrices. Fermentation produces beneficial effects in foods that undergo chemical changes caused by microorganisms such as bacteria or yeasts (Caplice and Fitzgerald 1999).

Fermentation plays different roles in food processing. Major roles considered are as follows:

- Preservation of food through formation of inhibitory metabolites such as organic acid (lactic acid, acetic acid, formic acid, and propionic acid), ethanol, bacteriocins and so on, often in combination with decrease of water activity (by drying or use of salt; Ross, Morgan and Hill 2002; Gaggia et al. 2011).
- Improving food safety through inhibition of pathogens (Adams and Mitchell 2002, Adams and Nicolaides 2008) or removal of toxic compounds (Hammes and Tichaczek 1994).
- Improving the nutritional value (Poutanen, Flander and Katina 2009, van Boekel et al. 2010).
- Organoleptic quality of the food (Marilley and Casey 2004; Smit, Smit and Engels 2005; Lacroix et al. 2010; Sicard and Legras 2011).

Biopreservation is a natural way to protect against spoilage and harmful contamination in food. This helps keeping food products fresh and safe throughout shelf life, opening for the possibility of reducing food waste. The food industry is presently looking for means of producing safe food products with an extended shelf life thus reducing food waste and meeting the consumer demands for natural, low salt, low sugar foods and for reduced use of chemical preservatives. Fermented food products have a longer shelf life and are less prone to spoilage than fresh food products of the same matrix (e.g. cheese compared to milk). There have been advances in the understanding of food microbiology, and the ability to screen for food cultures with better ability to stabilize food provides food cultures with bioprotective effect (Bech Hansen 2002). The advances of science in this field are eventually restricted by regulations in place, as this protective role of food cultures is mistakenly assimilated to the category of food additives and is not considered as a traditional use of food cultures in the food chain (Laulund et al. 2017). The aim of the present review is to highlight how the use of food cultures is already protecting the fermented food products, with enhanced shelf life and reduced spoilage.

The microbial stability and safety as well as the sensory and nutritive quality of foods are achieved by applying a combination of several different preservative factors called hurdles. The most

important hurdles for keeping food fresh and safe are temperature (high or low), water activity (a_w), acidity (pH), redox potential, preservatives (food additives), competitive microorganisms (bacteria, moulds and yeasts) and their metabolites. The competitive microorganisms with enhanced protective effect can be indigenous or be added as specifically selected food cultures.

As proposed by the International Dairy Federation on its fact-sheet on the topic (Available at: <https://fil-idf.org/publications/fr-ee-of-charge/idf-factsheet-007-2019-bioprotection/>) Biopreservation refers to enhanced food safety and extended shelf life of foods by indigenous and/or intentionally added microbiota, inhibiting growth of pathogenic and spoilage organisms due to microbiological competition and production of antimicrobial metabolites. Among the different species of food cultures, lactic acid bacteria have a major potential for use as biopreservation supported by their long history of safe use, proven antimicrobial properties, their capacity to naturally dominate the microflora and occupy the ecological niche during storage.

Food cultures for fermentation do all have biopreservation effects due to metabolically activities. While the traditional use of cultures in fermented foods refers to their positive action on product properties (texture, aroma, digestibility, ...), more specific focus is now on food cultures with appropriate biopreservative properties for a given application, inhibition of the spoilage microflora and improvement of food safety.

Biopreservation enhances the effectiveness of a food management system, but is never an alternative to good cleaning practices, hygienic design of the production and cold chain conservation (Motarjemi and Nout 1996). Food cultures with biopreservative effect create an extra protective hurdle against specific pathogens and/or spoilage microorganisms in the product during processing and/or after the product has left the manufacturing facility, e.g. during transport, storage, retail display and even after opening of the packaging by the consumer (Baka et al 2014).

The selection and application of protective food cultures must comply with the same safety criteria as for all food cultures used in the food industry. Food cultures are chosen for their ability to control and reduce foodborne pathogens and spoilage microorganisms by exploiting microbial competition and dominance phenomena. Isolation, selection, detailed characterization and validation of cultures are a way of taking advantage of the natural way in which microorganisms compete with each other in a complex environment in order to ensure that the added biopreservative food culture has the specific characteristics under specific condition(s). This allows more control over the process than with spontaneous fermentation. A spontaneous fermentation is actually also what takes place in fresh and ready to eat (RTE) foods resulting in spoilage. New analytical tools make it feasible to identify and characterize microorganisms

present in a given environment. These enable the selection of the best candidates from a very high number of food cultures to protect against pathogenic and spoilage microorganisms. Individual food culture strains within the same species have different fermentation properties that can create variations in taste, aromas and texture/viscosity that may or may not be wanted. A careful selection of individual strains can be based on well-known and commonly used species, but have to be tested individually regarding properties contributing to fit the purpose of certain biopreservative effects and possible side effects (Bourdichon et al. 2012).

GENETIC BASIS AND PRODUCTION OF SPECIFIC METABOLITES AND/OR ENZYMES

Microbes have the ability to produce an extraordinary array of metabolic by-products exhibiting a bactericidal or bacteriostatic activity such as organic acids, enzymes, bacteriocins and many other secondary metabolites. Biopreservation involves applying food-grade microorganisms to extend the shelf life of foods, and preventing the development of undesirable microorganisms (Elsser-Gravesen and Elsser-Gravesen 2014).

GENERAL AND LOW MOLECULAR WEIGHT METABOLITES

The end products of fermentation may include organic acids (lactate, acetate, succinate and formate), gases (CO_2 , H_2 and SO_2), hydrogen peroxide and other metabolites (ethanol, diacetyl, aldehydes, ketones, fatty acids and so on). The proportions of these end products vary between and within species. Many of these metabolites display antagonistic effects against other microbes through various modes of action (Ben Said et al. 2019). Examples include CO_2 which has long been documented as inhibitory to certain pathogenic species. Heterofermentative lactic acid bacteria (LAB) species produce CO_2 from formate, creating an anaerobic environment in the food which is hostile to aerobic species. Hydrogen peroxide (H_2O_2)-related inhibition occurs through oxidative damage of proteins and, at times, an increased membrane permeability take place in the target organisms. Another example is diacetyl (2,3-butanedione), produced by certain LAB species. While diacetyl can be employed as a biopreservative, its contribution to the flavor and aroma of a product means that it has limited preservative uses in foods where diacetyl is not wanted as a major flavor component. (Drosinos et al. 2005).

ANTIFUNGAL COMPOUNDS

Antifungal compounds are key to avoiding spoilage with yeasts and/or moulds, and a recent review describes the role and use of LAB as bioprotective food cultures against fungal spoilage in foods (Siedler, Balti and Neves 2019). Some compounds previously discussed, e.g. certain organic acids such as propionic acid, and reuterin can be used to inhibit fungi as well as bacteria (Corsetti et al. 1998, Cleusix et al. 2007). Phenolic acids produced by *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) are involved in the antifungal activity or in the reduction of aflatoxin production in e. g. maize (Nazareth et al. 2020). In other cases, antifungal compounds produced by LAB are peptides by chemical nature, such as those against *Aspergillus flavus* on maize (Muhialdin et al. 2020). Phenyllactic acid is another compound that can be used to inhibit both bacteria and fungi.

It is present in different forms such as 3-phenyllactic acid and 4-phenyllactic acid. Both are produced by LAB (Ström et al. 2002, Mu et al. 2010), whereas 3-phenyllactic acid can also be produced by *G. candidum* (Dieuleveux, Lemarinier and Guéguen 1998). In the latter case, the inhibition can also be towards *Fusarium* spp. and reducing the toxin concentration (Kawtharani et al. 2020). Since phenyllactic acid is produced by many LAB, it is often also naturally present in fermented foods and can be used in a variety of food products.

KILLER YEASTS

The multitude of fermented bacterial substances and mechanisms to control growth of other microorganisms, are similar known from yeasts fermentation. Killer yeasts produce and excrete extracellular proteins which are lethal to sensitive strains of other yeasts or other microorganisms (Starmer et al. 1987). These killer toxins (mycocins) were first detected in *Saccharomyces cerevisiae* and later in strains of other species such as *Debaromyces* spp., *Kluyveromyces* spp., *Pichia* spp., *Metschnikowia* spp. as well (reviewed in Mannazzu et al. 2019). Killer yeasts are applied as biocontrol agents especially in winemaking to control indigenous yeasts which produce offflavors (reviewed in Morata et al. 2020). Moreover, they have been used as bioprotectors in vegetables, olives, beer, sake, dry-cured ham, sausages, yogurt and cheese (reviewed in Salas et al. 2017, Medina-Córdova et al. 2018, Guimaraes et al. 2018, Mannazzu et al. 2019).

BACTERIOCINS

Bacteriocins are ribosomally synthesized peptides or proteins that possess antimicrobial activity towards closely related bacterial species, whereas the producing bacterium is immune to the specific bacteriocin itself (De Vuyst and Leroy 2007).

As examples, nisin has been chosen among lantibiotics (Class I bacteriocins), and lactococcin G and pediocin A among nonlantibiotics (Class II bacteriocins).

Nisin

Different strains of *Lactococcus lactis* subsp. *lactis* are known as bacteriocin producers. One of the most intensively studied and used is the lantibiotic nisin. Nisin is an effective agent against several undesirable Gram-positive bacteria in cheese and various other foods. It was also the first antimicrobial peptide approved by the FDA to be utilized as a food additive preservative (De Vuyst and Vandamme). The list of FDA approved food additives and GRAS (generally recognized as safe) notices with intended antimicrobial effect is provided in Table 1 (As available on December 2020–Available at: <https://www.fda.gov/food/gras-notice-inventory/recently-published-gras-notice-and-fda-letters>).

Nisin is a ribosomally synthesized lantibiotic and many variants have been discovered, naturally produced by several strains of *L. lactis* (i.e. Nisin A, Z, Q, F) and *S. uberis* (i.e. Nisin U, U2; Gross and Morell 1971; de Vos et al. 1993; Zendo et al. 2003; Wirawan et al. 2006; de Kwaadsteniet, Ten Doeschate and Dicks 2008). The most common variants are nisins A and Z, which differ in one amino acid residue. The biosynthesis and regulatory machineries of lantibiotics are encoded by genes organized in operons. Single genes can vary their location within the operon, but invariably they are clustered on the genome. For bacteriocin synthesis up to three functions are required: production, immunity and, optionally, quorum sensing. In the case of

Table 1. List of FDA approved food additives and GRAS (generally recognized as safe) notices with intended antimicrobial effect.

Food additives	Intended use	Food matrix
Acetic acid	Buffer and neutralizing agent	Cheese
Hydrogen peroxide	Used in combination with acetic acid to form peroxyacetic acid	Wash water for fruits and vegetable that are not raw agricultural commodities (59 ppm)
Lactic acid	Buffer and neutralizing agent	Bakery products, cheese, frozen desserts, fruit butters, jellies and preserves
Propionic acid	Preservative	Swiss and Gruyere Cheese
Natamycin (pimaricin)	Antimycotic	Cheese (< 20 mg/kg finished product)
Nisin preparation	Antimicrobial	Cheese (< 250 ppm)
GRAS-Microorganisms		
<i>Lactobacillus curvatus</i> DSM 18775	Antimicrobial (<i>Listeria monocytogenes</i>)	Ready-to-eat cooked meat, poultry products
<i>Lb. acidophilus</i> , <i>Lb. lactis</i> and <i>Pediococcus acidilactici</i>	Antimicrobial	Meat, poultry
<i>Carnobacterium maltaromaticum</i> CB1 (viable and heat-treated)	Antimicrobial (<i>Listeria monocytogenes</i>)	Various foods
<i>Carnobacterium maltaromaticum</i> CB1	Antimicrobial (<i>Listeria monocytogenes</i>)	Ready-to-eat meat products
GRAS-Bacteriocins		
Bacteriocin preparations specific to <i>Salmonella</i>	Antimicrobial (<i>Salmonella</i>)	Red meats, poultry and egg products (max. application rate 3 mg/kg or L)
Colicin preparations	Antimicrobial	Meat (application rate of 1–10 mg/kg)
Natamycin	Antimycotic (yeasts and molds)	Yogurt (levels < 5 ppm finished product)
Nisin	Antimicrobial	Casings of frankfurters; cooked meat; poultry products

nisin, these functions are scattered across the nisin biosynthesis gene cluster *nisABTCIPRKFE*G which consists of 11 genes divided into four operons (Lubelski et al. 2008). Bacteriocin production is encoded by *nisABTC* and *nisP*, *nisA* encoding the bacteriocin precursor, *nisT* the transporter exporting the unmodified precursor, *nisBC* encoding post-translational modification functions and *nisP* encoding the leader peptidase (Kuipers et al. 1993). Immunity is encoded by *nisI* and *nisEFG*, a dedicated immunity protein and an ABC-transporter, respectively (McAuliffe, Ross and Hill 2001). Lastly, the two-component signal transduction system responding to the mature nisin consisting of a response regulator and histidine kinase is encoded by *nisRK*, respectively (Ge et al. 2017). The self-protection mechanism of the food culture producers can involve more than one system. Comparative genomic analyses by Wels and collaborators (Wels et al. 2019) on publicly available genome sequences of *L. lactis* subsp. *lactis* and subsp. *cremoris* revealed a complete nisin biosynthesis cassette *nisABTCIPRKFE*G (Kuipers et al. 1993) on the chromosome of the subspecies *lactis* strains CV56 and IO-1. This gene cluster is flanked by transposase fragments. A complete nisin gene cluster was also found in the subspecies *lactis* strains KF134, KF146, KF196, KF282, K231, KF24, K337, KF67, KF7, Li-1 and LMG8526, at the same chromosomal insert position as in strain CV56. It was highlighted that all the strains of plant or vegetable origin can produce nisin Z. On the contrary, the subspecies *cremoris* strain V4 and the subspecies *lactis* strains LMG14418, LMG9446 and KF147 present an incomplete chromosomal gene cluster and cannot produce nisin, but they have maintained some immunity genes (i.e. *nisFEG* and/or *nisI*) (Wels et al. 2019). *Lactococcus lactis* ssp. *cremoris* FG2 and N41, from dairy starter and soil/grass, respectively, present a partial *nis* gene cluster encoding only *nisP*, *nisI* and a truncated *nisC*. This organization suggests a localization typical of plasmid (Tarazanova et al. 2016). The original nisin gene cluster is chromosome based. The detection of a plasmid based nisin variant produced by *Streptococcus capitis*

(O'Sullivan et al. 2020) and the detection of nisin variants in multiple species suggest that nisin like gene clusters can also be horizontally transferred on mobile elements.

Lactococcus lactis cheese starter cultures may either produce and/or tolerate the antimicrobial bacteriocin nisin. It is therefore relevant upon the choice of food cultures to assess their natural resistance or sensitivity to potentially produced bacteriocins (Van Gijtenbeek et al. 2021).

Pediocin PA-1

Pediocin, produced by *Pediococcus acidilactici*, belongs to the class II bacteriocins (nonantibiotics), a large and diverse group of antimicrobial compounds that includes small heat-stable, cationic and hydrophobic/amphiphilic peptides. They are mainly active against other LAB, and they damage target cells by pore formation or by interfering with the integrity of the membrane (Nes et al. 1996). Though most have a limited activity spectrum, some, including pediocin PA-1, also inhibit more distantly related bacteria. The key interest in pediocin PA-1 in relation to biopreservation is its functionality in inhibition of the pathogen *Listeria monocytogenes*. Where the nisin biosynthetic gene cluster is an example of a tightly controlled complex bacteriocin expression system, the pediocin PA-1 genetic organization features the other extreme, with a basic organization of a single gene cluster composed of only four genes *pedABCD* expressed in a single operon. Pediocin PA-1 production is ensured by the *pedA*, the pediocin precursor gene and *pedCD* encoding a dedicated ABC-transporter, while immunity is encoded by *pedB* (Rodriguez et al. 2002). Additional immunity systems as well as a quorum sensing system are absent. Interestingly, the cluster encoding pediocin PA-1 and highly similar bacteriocins is located on a plasmid that can be and likely has been transferred between species (Cui et al. 2012).

Lactococcin G

Lactococcins belonging also to the class II bacteriocins (nonlantibiotics). They are mainly active against other lactococci and they damage target cells by pore formation or by interfering with the integrity of the membrane (Nes et al. 1996). Of the two-peptide bacteriocins (class IIB), lactococcin G (LcnG) is the most studied in relation to its mode of action. It is constituted of the peptides LcnG- α (39 residues) and LcnG- β (35 residues) (Rogne et al. 2008), and its bactericidal activity relies on causing leakage of Na⁺ and K⁺ ions from the membrane of sensitive cells (Moll et al. 1996, 1998). The main target of lactococcin G is the membrane protein UppP/BacA, involved in the synthesis of peptidoglycan in the strains *L. lactis* ssp. *lactis* IL1403 and *L. lactis* ssp. *cremoris* MG1363 (Kjos et al. 2014).

Lactococcin G cluster is composed of two structural genes encoding the pre-bacteriocins (lagA and lagB), an immunity gene (lagC), an ABC transporter gene (lagD) and, located downstream of lagD, a gene coding for a transport accessory protein (lagE; Oppergard et al. 2010). The mechanism responsible for the secretion of lactococcin G by LagD and its dependency on the LagE transport accessory protein are not yet clarified.

Lactococcus lactis strains LMG2081 and BGBM50 are known as lactococcin G producers (Niessen et al. 1992; Mirkovic et al. 2015). The complete sequence of lactococcin G operon (~4.9 kb) in *L. lactis* LMG2081 has been deposited in the NCBI database (GenBank accession number FJ938036). BLAST alignment of the cluster against the database revealed the presence of a nearly complete LcnG operon (94% coverage, 97% identity) also on the chromosome of *L. lactis* strain CBA3619 isolated from kimchi.

OTHER ANTIBACTERIAL COMPOUNDS

The term bacteriocin-like inhibitory substances (BLIS) is used for presumptive bacteriocins still under investigation until their amino acid structure is identified (Settanni and Corsetti 2008). Other antibacterial substances fit into neither the low molecular weight metabolites, bacteriocins nor BLIS categories. They are not less active and sometimes have a broader spectrum of inhibition. Some of the compounds show limited inhibitory effect alone and are more active together with other substances or in combination with lactic acid (pH) producing food culture(s) (Niku-Paavola et al. 1999). This was demonstrated in cheese with a mixture of strains (Settanni et al. 2011), where not only antimicrobial activity was measured, but also increased growth of the starter cultures. Examples of such antimicrobial compounds showing a broad-range inhibitory activity include reuterin, a compound produced by *Limosilactobacillus reuteri* (formerly *Lactobacillus reuteri*) that inhibits fungi but also Gram-negative bacteria (Schaefer et al. 2010), and indeed, inhibition of *Clostridium difficile* has also been shown (Cleusix et al. 2007). A further example is D-3-Phenyl-lactic acid, shown to have inhibitory activity towards various pathogens like *Salmonella enterica*, *L. monocytogenes* and produced by several LAB but also *Geotrichum candidum* (Dieuleveux, Lemarinier and Guéguen 1998, Rodríguez, Martínez and Kok 2012).

Reuterin

As a bacterium occurring in sourdough, dairy and meat products, *Limosilactobacillus reuteri* (formerly *Lactobacillus reuteri*) has gained interest as a potential bioprotective food culture due to its ability to synthesize reuterin, a broad-spectrum antimicrobial system consisting of an isomeric mixture of

3-hydroxypropionaldehyde (3-HPA; Vollenweider et al. 2003). Reuterin displays inhibitory activity against bacteria, yeast, moulds and protozoa, including food spoilage and pathogenic organisms (Schaefer et al. 2010). Notably, metabolism of glycerol has been demonstrated to improve the competitiveness of *L. reuteri* in sourdough (Lin and Gänzle 2014). *Limosilactobacillus reuteri* uses a CoA-dependent pathway, in which 3-HPA is obtained from glycerol in a reaction catalysed by the coenzyme B12-dependent glycerol/diol dehydratase (GDH; Talarico and Dobrogosz 1990); 3-HPA is subsequently converted to 3-hydroxypropionic acid (3-HP) and 1,3-propanediol (1,3-PDO) (Dishisha et al. 2014). The glycerol/diol dehydratase of *L. reuteri* has been shown to be encoded by three genes located in the propanediol-utilization (pdu) operon (PduCDE; Morita et al. 2008); adjacent to this operon, *L. reuteri* possesses cbi-hem-cob genes that encode the proteins for the biosynthesis of vitamin B12 (Santos et al. 2008). In addition, Srimulu et al. (2008) found that glycerol/diol dehydratase is associated with microcompartments called metabolosomes, and their structural proteins are encoded by genes located in the pdu operon. The structure of the pdu-cbi-cob-hem cluster in *L. reuteri*, displaying a putative transposase gene between the pdu and cbi-cob-hem operons and IS elements within flanking regions, suggests this gene cluster may be a genomic island that has been acquired through horizontal gene transfer (Morita et al. 2008). Based on this hypothesis, it seems reasonable to assume that other species of LAB besides *L. reuteri* may have acquired this genetic island during evolution. Indeed, glycerol metabolism leading to 3-HPA production has been reported in *Secundilactobacillus collinoides* (formerly *Lactobacillus collinoides*; Sauvageot et al. 2000), *Loigolactobacillus coryniformis* (formerly *Lactobacillus coryniformis*) isolated from cheese (Martin et al. 2005), *Levilactobacillus brevis* (formerly *Lactobacillus brevis*) and *Lentilactobacillus buchneri* (formerly *Lactobacillus buchneri*; Schutz and Radler 1984), *Lentilactobacillus diolivorans* (formerly *Lactobacillus diolivorans*) from ciders (Garai-Ibabe et al. 2008). Consistent with this, the pdu operon has been detected in *L. collinoides* (Sauvageot et al. 2002) and in *L. brevis* (Makarova et al. 2006).

Phenyllactic acid

3-phenyllactic acid (2-hydroxy-3-phenylpropanoic acid, PLA) is acknowledged as a relevant contributor to the anti-microbial activity of LAB in fermented foods. A wide range of LAB genera, such as *Lactobacillus*, *Leuconostoc*, *Weissella*, *Pediococcus* and *Enterococcus* (Magnusson et al. 2003; Valerio et al. 2004; Ndagano et al. 2011; Li et al. 2014), have been demonstrated to produce PLA, though PLA biosynthesis has been most extensively studied in *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*; Lavermicocca et al. 2000; Ström et al. 2002; Prema et al. 2010; Wu et al. 2020). Remarkably, PLA exerts inhibitory effects *in vitro* and *in vivo* on several spoilage and mycotoxigenic moulds from sourdoughs and bakery products (Lavermicocca, Valerio and Visconti 2003; Dal Bello et al. 2007; Ryan et al. 2011; Valerio et al. 2016).

In LAB, PLA is a by-product of phenylalanine (Phe) catabolism: phenylalanine is firstly transaminated to phenylpyruvic acid (PPA) by an aromatic aminotransferase (AAT; Yvon et al. 1997) and subsequently reduced to PLA by a 2-hydroxyacid dehydrogenase (2-HADH) such as lactate dehydrogenase (LDH; Vermeulen, Ganzle and Vogel 2006; Li, Jiang and Pan 2007; Mu et al. 2010). Although the genes encoding such enzymes are ubiquitously present in LAB, significant differences are recorded in the amount of PLA produced by different strains, and this disparity was ascribed to varying enzymatic

activity of LDH toward PPA (Li et al. 2008). Recently, it has been demonstrated that *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) LY-78 synthesizes PLA de novo via the Phe synthetic pathway and suggesting that panE1 (ketopantoate reductase), serA1 (D-3-phosphoglycerate dehydrogenase) and ldhD2 (D-lactate dehydrogenase 2) may be key genes of PLA biosynthesis in LAB (Sun et al. 2019).

CONTROL OF ACIDITY OF FERMENTED FOOD PRODUCTS AS A RESULT OF FERMENTATION

Acidification mechanism

Many parameters govern the survival and growth of microorganisms in food. The acidity or pH of a food can affect the type and number of microorganisms present. All microorganisms have an optimum pH value for growth, and altering the hydrogen ion concentration can influence the growth of an organism or even inhibit growth. In general, bacteria prefer to grow at a pH near neutrality (pH 6.5–7.5) but will tolerate a pH range of 4–9. Yeasts are more tolerant of lower pH values than bacteria, while moulds survive across the widest range of pH values. Foods with a pH value below 3.5 can support the growth of both yeasts and moulds. Because of the sensitivity of organisms to widely differing pH values, the pH provides a powerful selection which influences the species or group of microorganisms that will predominate in unaltered food products. For example, bacteria primarily spoil proteinaceous foods such as dairy, meat, poultry and seafood with a pH range of 5.5–6.5. In contrast, yeasts and moulds more commonly proliferate on fruits and vegetables with inherently lower pH values and little buffering capacity (Doores 2005).

One effective means to preserve food from spoilage is to increase the acidity of the food thereby creating an unfavourable environment for survival of undesirable microorganisms through natural fermentation. Depending on the final pH, this inhibition can be either biostatic or biocidal. The final result will depend on the microbial species, the type and concentration of the acidulant, the time of exposure, the buffering capacity of the food and most likely the compositional/chemical properties of the food (Doores 2005).

Microorganisms display varied sensitivity to acids. In most cases, they are vulnerable to the organic acids they produce because they are by-products of primary metabolism and as such are their natural 'electron sink'. It may happen, however, that in mixed culture fermentations, the produced acid is a source of energy for another microorganism. For example, LAB produce lactic acid as a by-product of their metabolism, which serves as an energy source for propionic acid bacteria (PAB) and the mould *Penicillium roqueforti*. *St. thermophilus* produces formic acid as a by-product of their metabolism, which serves as an energy source for *L. bulgaricus* in yogurt fermentation.

Organic acids are weak acids, so they do not dissociate completely in aqueous environments, and their antimicrobial activity depends on the degree of dissociation and the pH in the food environment. Hence, the antibacterial activity of organic acids is increased when the pH of the food is low. A decrease in pH leads to an increase in protonated acid concentration, decreasing the polarity of the molecules and consequently increasing the diffusion of acids across the cell membrane and into the cytoplasm (Doores 2005; Mani-López, García and López-Malo 2012). The inhibitory effect of organic acids is based on pH, concentration, chain length, type and degree of branching. Indeed, effective use of an acidulant depends on the dissociation constant

(pKa) or the pH at which 50% of the total acid is dissociated. The pKa of most organic acids is between pH 3 and 5. The pKa of the acids most commonly found in fermented foods are: acetic acid, pKa 4.75; lactic acid, 3.08 and propionic acid, 4.87. Because the undissociated portion of the molecule is believed to be responsible for the antimicrobial effect, it would be advantageous to use the acids near these values from a biopreservation perspective. As the pH of a solution decreases, the concentration of the undissociated form will increase for all acids. For weaker acids, the undissociated proportion at any given pH will be higher than for stronger acids, i.e. those with a lower pKa (Adams 2014).

In foods having a range of buffering components such as proteins and amino acids, it is not easy (possible) to calculate a degree of dissociation of weak acid using the Henderson-Hasselbach equation. The issue may be further complicated by the presence of oil or fat into which the acid might partition preferentially. This would have the effect of decreasing the acidity in the aqueous phase in which microbial growth occurs and thereby the anticipated antimicrobial effect (Wilson, Wilson and Waspe 2000).

Because pH values below 4.5 will stop or severely curtail the growth of all the major bacterial pathogens and will, depending on the conditions, ultimately lead to their death/inactivation, food safety concerns tend to be much reduced when considering acidic/fermented foods. It has long been recognized that *Clostridium botulinum* spores will not germinate and grow at pH values below 4.6, and this is enshrined in various codes of practice. Usually, this organism can be well controlled by efficient acid production. Levels of acidity which do not kill pathogens or stop their growth entirely can still improve food safety. The risk from infectious pathogens such as *Salmonella* spp. will be lower if growth and thereby numbers of the organisms are restricted and, at suboptimal pH, toxigenic organisms such as *Staphylococcus aureus* cannot grow to levels sufficient to produce biologically effective concentrations of toxin in the food. This is exemplified in EU regulations where food safety criteria for milk powder and some cheeses specify that only when levels of coagulase positive staphylococci exceed 10^5 CFU/g, there is a requirement to test for enterotoxin content (Adams 2014).

Lactic acid fermentation

Lactic acid is produced by many microorganisms of which the most known are LAB belonging to genera former *Lactobacillus* genus, *Leuconostoc*, *Pediococcus*, *Lactococcus* and *Streptococcus* (Crowley, Mahony and van Sinderen 2013).

There are three different pathways leading to production of lactic acid from glucose (lactose); substrate level phosphorylation (homofermentative process) leading to production of two molecules of lactic acid from one of glucose; 6P-Gluconate pathway (heterofermentative process) leading to production of one molecule of lactate, ethanol and carbon dioxide from one molecule of glucose; and the Bifidus pathway leading to production of two molecules of lactate and three acetate from two molecules of glucose (Kandler 1983). The produced acid is in the form of L(+), D(-) or as racemic lactic acid D(-) and L(+). The production and secretion of lactic acid and other weak organic acids results in an acidic environment which generally restricts growth of both bacteria and fungi, including many pathogenic and spoilage microbes (Ross, Morgan and Hill 2002). Acid production and a simultaneous reduction in pH are inevitable consequences of LAB growth, and acidity levels in some fermentations can exceed 100 mM, reducing the pH to below 4.0 in weakly buffered systems (Adams and Mitchell 2002).

The rate of pH drops and the final pH value in lactic acid fermentations depend on a number of factors such as the buffering capacity and water activity of the medium, the temperature and duration of fermentation, the inoculum size and the metabolic activity of the bacteria. Ideally, the target pH would be around 4.5, although this is not achieved in many common fermented foods such as cheese. Even in very weakly buffered media, the pH would tend to level off around 3.8 as lactic acid production produces a lactate buffer. Maximum effect will also be achieved if the pH drop occurs rapidly, within hours, to prevent any pathogen growth occurring (Adams 2014).

Acetic acid fermentation

The microorganisms oxidizing ethanol to acetic acid are commonly called acetic acid bacteria (AAB). Acetic acid bacteria are mesophilic obligate aerobes that oxidize sugars, sugar alcohols and ethanol, with the production of acetic acid as the major end-product. During acetic acid production, ethanol is almost quantitatively oxidized to acetic acid. Acetic acid bacteria exhibit resistance to high acetic acid concentrations and low pH (Raspor and Goranović 2008; Yamada and Yukphan 2008). Physiologically, bacteria belonging to the genus *Acetobacter* sp. convert alcohols to acids by oxidation.

Propionic acid fermentation

Propionic acid bacteria (PAB) are combined into the family *Propionibacteriaceae*, genus *Propionibacterium* and *Acidipropionibacterium*. They are isolated from milk, fermented dairy products and cheese; they are also found in at least 24 different vegetables and fruits species (Vorobjeva, Khodjaev and Vorobjeva 2008).

Depending on the strains, the ratio of PA to AA can vary widely and well beyond theoretical 3:1. Their product, PA alone or with AA, is inhibitory toward *Aspergillus flavus*, aerobic *Bacillus*, *Salmonella* spp. and yeasts and has been used as a mould inhibitor for animal feed, wet corn, silage and grain (Balamurugan, Venkata and Panda 1999) as well as in the food industry to prevent spoilage of foods such as bread and cake from moulding.

ECOLOGICAL COMPETITION

Microorganisms inhabit almost every environment in the world, including different food matrices. Interactions between different microorganisms are unavoidable and can either be symbiotic or competitive. Mechanisms of interaction include e.g. the production of inhibitory molecules and the competition for nutrients. Andreevskaya et al. (2018) showed that *Leuconostoc gelidum*, a spoilage LAB, in packaged cold-stored meat products enhanced its nutrient-scavenging capabilities in the presence of *Lactococcus piscium* and *Paucilactobacillus oligofermentans* by upregulation of carbohydrate catabolic pathways, pyruvate fermentation enzymes and ribosomal proteins. The slower growing *Lc. piscium* and *P. oligofermentans* downregulated these functions in the presence of *Le. gelidum*, but overexpressed prophage genes and restriction modification systems, which are mechanisms of DNA exchange and protection against it (Andreevskaya et al. 2018). Several studies on growth competition of spoilage organisms and potential bioprotective strains in media or food models exist. The mechanism of action responsible for growth inhibition is unfortunately often not examined. As an example: Leyva Salas et al. (2018) tested the antifungal activity of 32 LAB

strains against the four fungi *Penicillium commune*, *Mucor racemosus*, *Galactomyces geotrichum* and *Yarrowia lipolytica* to find a combination of food cultures, which has a bioprotective effect without influencing the organoleptic properties nor inhibiting starter cultures of the respective foods. They used a cheese and yogurt model system and received different outcomes for the respective cultures in the different models, showing that food matrix and production conditions play a significant role in determining antifungal activity of potential bioprotective food cultures.

A way for microorganisms to colonize a favourable environment is either to colonize this environment as soon as it is available, e.g. through fast growth, or to actively displace already existing microorganisms, e.g. by production of antimicrobial compounds (Hibbing et al. 2010). When considering antifungal activity of LAB, many studies have revealed that antifungal compounds are often found in concentrations below the minimal inhibitory concentration (MIC), thus pointing out that other mechanisms must be responsible for the bioprotective effects of LAB against yeasts and moulds as well (Siedler, Balti and Neves 2019). The competition for limited resources, for example nutrients, carbon source and essential ions, can result in inhibition or delay of growth of either protective cultures or spoilage organisms. Honoré et al. (2016) found that the inhibition of *Penicillium* spp. by *Lacticaseibacillus paracasei* (formerly *Lactobacillus paracasei*) in a defined medium was not only induced by metabolite formation, but also by nutrient consumption, especially by the consumption of glucose and glutamine. Reduced glucose availability induced by *Penicillium chrysogenum* growth was also assumed to be the reason for repressed Ochratoxin A (OTA) production of *Penicillium nordicum* (Delgado et al. 2019).

Furthermore, nitrogen is a limiting factor for microbial growth in dairy-based systems. Micro-organisms initially compete for free amino acids and small peptides, while they compete for peptides in later stages of fermentation. The ability to utilize amino acids efficiently is essential for growth of the respective microorganism (reviewed in Sieuwerts et al. 2008). Transcriptome analysis was used to study the interactions of *Lc. lactis* and *Saccharomyces cerevisiae* in coculture during the exponential growth phase. Whereas the lactic acid concentration was the same in a single culture of *Lc. lactis* and in the mixed culture of *Lc. lactis* and *S. cerevisiae*, ethanol concentration and glucose consumption were increased in the mixed culture. Also, pyrimidine metabolism of *Lc. lactis* was reoriented, most likely regulated by the ethanol production of the yeast (Maligoy et al. 2008).

Manganese was found to be the limiting factor for dairy yeast and mould growth in yogurt with *Lacticaseibacillus rhamnosus* (formerly *Lactobacillus rhamnosus*) and *Lb. paracasei* as protective food cultures (Siedler et al. 2020). The manganese transporter (MntH1) is responsible for manganese uptake under acidic conditions and gives strains expressing the *mntH1* gene the ability to take up manganese. The resulting manganese depletion ends up in delayed yeast and mould growth. Homologues of the *mntH1* gene were found in 15 different *Lactobacillus* species, indicating that this mechanism of competitive exclusion could be a general mechanism of LAB to interfere with yeast and mould growth (Siedler et al. 2020).

Iron is another nutrient which is fundamental to bacterial growth and thus a benefit for some micro-organisms if they are able to produce iron scavenging molecules, so called siderophores, to acquire iron from the environment (Hibbing et al. 2010). This has been shown for ripening bacteria on the surface of smear-ripened cheese. The addition of either iron or the

siderophore desferrioxamine B stimulated growth of *Arthrobacter* spp., *Corynebacterium* spp. and *Brevibacterium* spp. Furthermore, genes for iron-siderophore transporter binding proteins of *Arthrobacter arilaitensis* were upregulated when the siderophore was added, but not when iron was added to the medium (Monnet, Back and Irlinger 2012). Sipiczki (2006) hypothesized that contrary to bacterial siderophore production, *Metschnikowia* spp. inhibits growth of other micro-organisms by immobilizing iron in the medium due to formation of an insoluble pigment. Additionally, iron depletion by the biocontrol yeast *Metschnikowia pulcherrima* was successfully used to control the postharvest apple-pathogens *Botrytis cinerea* and *Alternaria alternata* during apple storage (Saravanakumar et al. 2008).

As soon as a microorganism encounters a favourable environment, binding and attachment are crucial to colonize the ecological niche. Many pathogenic bacteria are secondary colonizers of biofilms (Giaouris et al. 2015). Shaping the biofilm present on food (e.g. smear cheese) or processing environment (e.g. stainless steel) through the application of protective cultures is a challenging but promising approach. As an example: Habimana et al. (2009) studied the attachment of *L. monocytogenes* on biofilms formed by different customized *L. lactis* strains and found that the adhesion of planktonic cells was almost prevented when exopolysaccharides were formed by the biofilm-forming cells, but the adhesion was increased when biofilms had a porous structure formed by chain-making strains.

FOOD CULTURES AND BIOPRESERVATION: INDUSTRIAL APPLICATIONS

In ancient times, food fermentation was a spontaneous event whose outcome was uncertain in terms of quality and safety. Nevertheless, it was a first step for improving the shelf life (Farnworth 2008). Through the centuries, the monitoring of this process regularly improved, from empirically developed good practices in households to a systematic documentation of HACCP procedures in modern food processing industry. Next to pasteurization, salting and other hurdles, the application of food cultures in traditional fermented foods such as yoghurt, cheese, fermented meat, vegetables, beer and wine is nowadays strongly established. There is also a trend towards the use of traditional biotechnology for developing new foods and beverages (Laranjo, Potes and Elias 2019). Food cultures produce a high variety of compounds, including organic acids, alcohol and aroma compounds, contributing to product texture, taste and safety. This biopreservation contribution to safety meets the growing demand of the consumers for minimally processed food products.

The application of food cultures, which is widespread in dairy, meat and vegetable products, is already a form of biopreservation, but the addition of protective food cultures with a better productive effect can add another hurdle against pathogens or spoilage organisms. This is especially the case for fresh and RTE food products which lack a heating treatment and contain therefore potential spoilage organisms.

Applied research in the area of protective effect of food cultures is a challenging task. Between the discovery of an inhibitory activity in synthetic media and a commercially available food culture, a protective food culture has to pass the so-called challenge tests. It is thereby inoculated in a given food matrix together with the target microorganism and evaluated for the ability to reduce or control the outgrowth of the target organism over the desired shelf life. Applied research has been

carried out extensively in the last two decades focusing on inhibition of pathogenic microorganisms such as *L. monocytogenes*, *Salmonella* sp., Shigatoxin producing *Escherichia coli* and *Staphylococcus aureus* as well as spoilage microorganisms such as yeasts, moulds and *Clostridium* species (Settani et al. 2008; Castellano et al. 2017; Leyva Salas et al. 2017; Bosse Née Danz et al. 2018; Oliveira et al. 2018; Silva, Silva and Ribeiro 2018; Laranjo, Potes and Elias 2019; Rouse and van Sinderen 2008). Well-adapted microorganisms will provide a more efficient protection with more effective inhibitory microorganisms often being isolated from the investigated food itself (Austin-Watson, Grant and Brice 2013; Lee et al. 2016; Scatasa et al. 2017). However, care must be taken that those cultures do not have a negative impact on the organoleptic characteristics of the end-product.

Studies covering a wide range of foods including all fermented foods listed above as well as non-fermented foods are presented below.

Dairy products

As first line of defence, fermented dairy products are preserved through the acidification carried out by food cultures, as illustrated by the high safety of yoghurt. Thanks to its low pH, yoghurt is only susceptible to yeast and moulds (Leyva Sala et al. 2017), but recently a general mechanism of LAB to also inhibit yeast and mould growth was discovered (Siedler et al. 2020). The careful selection of strains constituting mesophilic food culture of mildly acidified products such as cheese can already substantially increase the protection against pathogens and spoilage agents, as extensively shown for the species *L. lactis* (Silva, Silva and Ribeiro 2018). The applications mainly cover the protection against *L. monocytogenes* and *Clostridium tyrobutyricum* (Garde et al. 2011; Kondrotiene et al. 2018; Lianou and Samelis 2014; Samelis and Kakouri 2018). *Lactobacillus* spp. and *Enterococcus* spp. as an adjunct culture may also increase food safety (Cocolin et al. 2007; Martinez et al. 2015). Raw milk soft cheeses are highly sensitive to contamination by gram-negative bacteria such as *Salmonella* spp. and Shigatoxin producing *E. coli*. Application of LAB together with the gram-negative species *Hafnia alvei* was shown to protect those cheese types (Callon, Arliguie and Montel 2016). Non-fermented dairy products such as cottage cheese also benefit from the addition of selected protective food cultures as shown by numerous studies (Silva, Silva and Ribeiro 2018; Chhetri, Prakitchaiwattana and Settachaimongkon 2019).

By analogy to bioprotection observed in traditional fermented food, a combination of strains is often more powerful than a single strain (Aljasir et al. 2020; Aljasir and D'Amico 2020; Rodriguez et al. 2012; Chhetri, Prakitchaiwattana and Settachaimongkon 2019; Sindi et al. 2020).

While research mainly focuses on LAB, further microorganisms isolated from the food microbiota may also contribute to food safety. One research field focuses on the rind of ripening cheeses that may support survival of pathogens (Roth et al. 2011; Imran et al. 2013; Callon et al. 2014). As the biodiversity of rind microbiota is by far higher than the core microbiota, the development of future protective food cultures relies on a deep understanding of the species interactions in this habitat. In many cases, such added food cultures produce inhibitory compounds against various pathogens or spoilage bacteria, but in certain cases a combined effect of the starter together with the protective food culture inhibits growth. In many cases, *L. lactis* subsp. *lactis* is involved either as protective food culture or as starter culture (Kondrotiene et al. 2018), but other species

such as *Latilactobacillus sakei* (formerly *Lactobacillus sakei*) or *Enterococcus* spp. can also be used and this demonstrates that no regulatory difference should be made (Cocolin et al. 2007; Martinez et al. 2015). Another application is the use against *Staphylococcus aureus* (Aljasir and D'Amico 2020). In certain cheeses such as cottage cheese, halophilic food cultures can be used also against *S. aureus* (Chhetri, Prakitchaiwattana and Settachai-mongkon 2019).

The prevention of growth of *Salmonella* spp. in dairy products can also be achieved using specific cultures added to dairy products. *Salmonella* spp. was inhibited in cheese using a *Hafnia alvei* food culture, and the same authors showed the increased inhibitory effect of the combination of selected food cultures against *L. monocytogenes* (Callon, Arliguie and Montel 2016; Aljasir et al. 2020).

A combination of several species/strains for the inhibition of pathogens, which was usually the case in the early days of fermentation, can enhance the antimicrobial activity due to either increased inhibitory power of a wider range of target organisms. This is, for example, the case for kefir products which has been described to have antimicrobial properties attributed to its low pH and specific antimicrobial substances produced during the fermentation process (Kim et al. 2016; Sindi et al. 2020).

Meat and fish products

There are various hurdles used to preserve meat and fish products, including addition of salt, nitrite, starter cultures, smoking and cold storage. There is increasing evidence that protective food cultures may also play a significant role against *L. monocytogenes*, toxin-producing *Staphylococcus aureus* and further pathogenic and spoilage microorganisms (Castellano et al. 2017; Bosse Née Danz et al. 2018; Oliveira et al. 2018; Laranjo, Potes and Elias 2019; Aljasir et al. 2020). In meat, nitrite is a salt with potential toxic effects, and there is pressure to reduce its use. A combination of nitrite with a protective food culture demonstrated a higher reduction of *L. monocytogenes* in fermented sausages than without the addition of the protective food culture (Nikodinoska et al. 2019). The combined effect of food cultures has been shown to inhibit Ochratoxin A production in dry cured ham by *Penicillium chrysogenum* and *Debaryomyces hansenii* (Cebrian et al. 2019).

Since fish and meat are not sterile (just like RTE Food products), their own microbiota will be active during storage. Psychrophilic microbiota already present on meat and fish such as *Carnobacterium* spp. and *Lactococcus piscium* can inhibit growth of pathogenic and spoilage bacteria. This is very promising, but careful selection of strains has to be carried out, as those highly adapted species will either spoil or protect the product, depending on their strain-specific phenotypes. (Castellano et al. 2017; Zhang, Gänzle and Yang 2019; Bazarnova et al. 2020)

Vegetables and cereal products

Moulds are the major spoilage issue when it comes to preservation of vegetables and cereal products. For a detailed overview of antifungal protective cultures in this type of food, we suggest reading the following recent review (Leyva Salas et al. 2017).

Listeria monocytogenes is also a pathogen of concern for vegetables. The application of protective cultures can be either directly on the product or through the washing process. Ramos et al. (2020) showed that a *Pediococcus pentosaceus*, prevented *Listeria* sp. proliferation in vegetable. In cabbage, the application of a *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) strain reduced *L. monocytogenes* but showed at the same time

that it is also important to synchronize all hurdles since the absence of oxygen increased the resistance of *L. monocytogenes* against *L. plantarum* (Dong et al. 2020).

Sourdough bread is a fermented product using yeasts and LAB. This microbiota can naturally protect the product from spoilage such as moulds (Chavan and Chavan 2011). In the case where the fermentation process for bread is mainly dominated by yeast, the addition of specific LAB food cultures enables the inhibition of for example *Aspergillus* spp., *Penicillium* spp. and *Fusarium culmorum* (Russo et al. 2017), thereby increasing safety (reduction of mycotoxins) and quality (reduction of off-flavour).

Industrial protective food cultures

Thanks to extensive scientific investigation in the last decades, food producers start to have access to protective cultures specifically developed to control a given pathogenic or spoilage microorganism in a given food. Compared to the tremendous number of studies dedicated to the development of protective food cultures illustrated above, there are still only a few available on the market. One cause may be that the protective food culture has to fit into an already intricate hurdle concept. Therefore, models used in future investigations should be as close as possible to the real processing conditions. In particular, care must be taken for commercial bioprotective food cultures not to have a negative impact on the organoleptic characteristics of the product. The regulation of use and declaration of protective cultures nevertheless remains an intricate topic for regulators in some regions/countries. The protective and preservation effects are seen as a key new effect of food cultures and it is regarded as belonging to food additive.

SAFETY DEMONSTRATION OF FOOD CULTURES FOR THEIR BIOPROTECTION ACTIVITIES

There is currently no firmly established regulation for the safety assessment of live micro-organisms added to food products as cultures or ingredients. However, there are a number of guidelines, recommendations and expert reviews on possible steps to document and validate the safety of live microorganisms used in foods independent of the mode of action of the food cultures (Laulund et al. 2017). Most industrial food strains used today are bacteria from species with a history of use in food products without apparent adverse effects. Four types of investigations have been proposed as further detailed.

Opportunistic infections

Commensal bacteria have been described to cause infections in patients with underlying disease (Berg and Garlington 1979; Berg 1985,1995). Owing to its natural presence in different sites of the human body and in fermented food products, the genus *Lactobacillus* has gained particular attention. *Lactobacillus* infections occur at a very low rate in the generally healthy population – estimated 0.5/1 million per year (Borriello et al. 2003; Bernardeau, Guguen and Vernoux 2006). As stated in two reviews of *Lactobacillus* infections: 'Underlying disease or immunosuppression are common features in these cases, whereas infection in previously healthy humans is extremely rare' (Aguirre and Collins 1993), and 'Lactobacillus bacteraemia is rarely fatal per se but serves as an important marker of serious underlying disease' (Husni

et al. 1997) sporadic infections have been reported in immunocompromised patients. The underlying problems have mainly been central venous catheter (CVC) in place, metabolic disorders, organ failure, or invasive procedures as dental work (Axelrod et al. 1973; Liong 2008). Infections by other bacterial species used as food cultures are also extremely rare (Horowitz et al. 1987; Barton, Rider and Coen 2001; Mofredj, Bahloul and Chanut 2007; Leuschner et al. 2010).

Infections with the commonly used yeast and mould species are rare events as well (Enache-Angoulvant and Hennequin 2005). Most of the infections are due to opportunistic pathogens not recognized as food cultures and affect immunocompromised patients and hospitalized patients (Jacques and Casaregola 2008; Miceli, Diaz and Lee 2011). In the 2018 re-evaluation EFSA concluded: "The safety concerns described are all considered linked to severe underlying health conditions and therefore do not change the consideration of *Lactobacillus* spp. for the QPS status" (EFSA 2018).

Toxic metabolites and virulence factors

Biogenic amine formation in fermented foods by LAB has recently been reviewed (Spano et al. 2010). Following food poisoning outbreaks (Sumner et al. 1985), metabolic pathways have been elucidated (Straub et al. 1995) and screening procedures proposed to limit the level of production (Bover-Cid and Holzapfel 1999; Bover-Cid, Izquierdo-Pulido and Vidal-Carou 2000).

The presence of mycotoxin genes also raises safety concerns, although the level of expression within fermented food is very unlikely to cause any health hazard (Barbesgaard, Heldt-Hansen and Diderichsen 1992). Within fungi, the potential for antibiotic production is also an undesired property.

The occurrence of virulence traits should not be present in microorganisms used in a food fermentation. A specific risk assessment should be conducted on strains presenting these undesirable properties, even if they belong to a species with a long history of use (Semedo et al. 2003a, b).

Antibiotic resistance

The emergence and spread of antibiotic resistance are a major global health concern. The on-going Codex ad hoc intergovernmental task force on antimicrobial resistance is focused on the non-human use of antimicrobials. Microorganisms intentionally added to food and feed for technological purposes have not been shown to aggravate the problem of spreading antibiotic resistant pathogens (Anonymous 2011).

Intrinsic resistance or resistance that is caused by mutation in an indigenous gene not associated with mobile elements would represent a very low risk of dissemination (Saarela et al. 2007). Acquired antibiotic resistance genes, especially when associated with mobile genetic elements (plasmids, transposons), can be transferred to pathogens or other commensals along the food chain from within the product until consumption (FEEDAP 2005, 2008; Nawaz et al. 2011).

The role of microorganisms in the spread of antibiotic resistance has been assessed in fermented foods (Nawaz et al. 2011). Results of such studies confirm the role of a reservoir of antibiotic resistance genes from the food microbiota, without identifying any major health concerns to date.

It is considered that strains carrying acquired antibiotic resistance genes might act as a reservoir of transmissible antimicrobial resistance determinants (FEEDAP 2005, 2008).

Gene transfer of antibiotic resistance between microorganisms in the food and feed chain is thus considered to be a topic of surveillance for the safety demonstration of microorganisms (Borriello et al. 2003; Gueimonde et al. 2005).

Definition of 'History of use'

The concept of 'history of safe use' has appeared recently in regulations and in safety assessment guidance. One definition of 'history of safe use' proposes "significant human consumption of food over several generations and in a large, genetically diverse population for which there exist adequate toxicological and allergenicity data to provide reasonable certainty that no harm will result from consumption of the food" (Health Canada 2003). In order to evaluate the history of safe use of a microorganism, it is necessary to document not just the occurrence of a microorganism in a fermented food product, but also to provide evidence of whether the presence of the microorganism is beneficial, fortuitous, or undesired.

CONCLUSION

Biopreservation of a food product can be achieved by fermentation, in a targeted or untargeted way. While the different bacteriocins described (mostly from LAB) show a dedicated mechanism to ensure primarily the survival in an ecosystem of a microbial species, but serve also as an additional hurdle to ensure safe food, other metabolic pathways, such as acidification, most specifically preserve initially short life stable food matrices (uninoculated milk get spoiled within hours). The initial empiric use of fermentation in the food chain was done for extended shelf life and avoiding food waste and food lost. Adaptation to texture and sensorial properties came after. The consumer has adapted its taste preferences to the different food cultures, and what is fermented food for one population and part of its culture can be considered spoiled if not dangerous by others.

Biopreservation itself can be ensured by other biological mechanisms than fermentation that were not considered in the present review. They would require a different approach, both for usage and safety demonstration. The potential of application of food cultures for biopreservation should not be considered something 'new' *per se* requiring a specific regulation and application process. It remains a traditional use of food cultures in the food chain, most presumably its first use. The focus in the initial steps of food microbiology on the sensorial properties has had a counterproductive effect on the application of food cultures for other applications. Fermentation applied to biopreservation should not be understood as a new approach, but one that time made us forget. Its potential as an additional hurdle to tackle food waste in combination with the numerous control measures already in place should not be underestimated. Enabling the delivering of safe, stable and tasty foods fitting in a sustainable lifestyle.

ACKNOWLEDGMENTS

Although the present review is not endorsed by the International Dairy Federation (IDF), the authors, most of them delegate of their respective countries to the Standing Committee on Microbiological Hygiene (SCMH), are also members of an IDF action team on Fermentation and biopreservation. They want to thank the head office and the national committees for their support on this project that will led to a publication to clarify the link between fermentation and bioprotection through the action of

Food Cultures to address eventual concern regarding the safe use of microorganisms in fermented foods.

Conflicts of Interest. None declared.

REFERENCES

- Adams M, Mitchell R. Fermentation and pathogen control: a risk assessment approach. *Int J Food Microbiol* 2002;**79**:75–83.
- Adams MR, Nicolaides L. Review of the sensitivity of different foodborne pathogens to fermentation. *Food Control* 2008;**8**:227–39.
- Adams MR. Acids and Fermentation. In *Food Safety Management – A Practical Guide for the Food Industry* ed. Yasmine Motarjemi and Huub Lelieveld, Elsevier, 2014, 467–79. DOI: 10.1016/B978-0-12-381504-0.00019-6.
- Aguirre M, Collins MD. Lactic acid bacteria and human clinical infection. *J Appl Bacteriol* 1993;**75**:95–107.
- Aljasir SF, D'Amico DJ. The effect of protective cultures on *Staphylococcus aureus* growth and enterotoxin production. *Food Microbiol* 2020;**91**:103541. DOI: 10.1016/j.fm.2020.103541.
- Aljasir SF, Gensler C, Sun L et al. The efficacy of individual and combined commercial protective cultures against *Listeria monocytogenes*, *Salmonella*, O157 and non-O157 shiga toxin-producing *Escherichia coli* in growth medium and raw milk. *Food Control* 2020;**109**:106924. DOI: 10.1016/j.foodcont.2019.106924.
- Andreevskaya M, Jääskeläinen E, Johansson P et al. Food spoilage-associated *Leuconostoc*, *Lactococcus*, and *Lactobacillus* species display different survival strategies in response to competition. *Appl Environ Microbiol* 2018;**84**:e00554–18.
- Anonymous. Codex Alimentarius Guidelines for Risk Analysis of Foodborne Antimicrobial Resistance. CAC/GL 77–2011, 2011, 1–29. Available at: (August 2020) www.fao.org/input/download/standards/11776/CXG.077e.pdf.
- Austin-Watson C, Grant A, Brice M. Suppression of *Listeria monocytogenes* by the Native Micro-Flora in Teewurst Sausage. *Foods* 2013;**2**:478–87.
- Axelrod J, Keusch GT, Bottone E et al. Endocarditis caused by *Lactobacillus plantarum*. *Annales Internes de Médecine* 1973;**78**:33–7.
- Baka M, Fernández EN, Mertens L et al. Protective role of indigenous *Leuconostoc carnosum* against *Listeria monocytogenes* on vacuum-packed Frankfurter sausages at suboptimal temperatures. *Food Res Int* 2014;**66**:197–206.
- Balamurugan K, Venkata DV, Panda T. Propionic acid production by whole cells of *Propionibacterium freudenreichii*. *Bioprocess Eng* 1999;**20**:109–16.
- Barbesgaard P, Heldt-Hansen HP, Diderichsen B. On the safety of *Aspergillus oryzae*: a review. *Appl Microbiol Biotechnol* 1992;**36**:569–72.
- Barron FH, Fraser AM. Acidified foods: food safety considerations for food processors. In: Muzzalupo Innocenzo (ed). *Food Industry* 2012, 231–9.
- Barton LL, Rider ED, Coen RW. Bacteremic infection with *Pediococcus*: vancomycin-resistant opportunist. *Pediatrics* 2001;**107**:775–6.
- Bazarnova J, Korableva N, Ozerova O et al. Biochemical composition and quality of herring preserves with addition of bio-protective cultures. *Agron Res* 2020;**18**:1629–39.
- Bech Hansen E. Commercial bacterial starter cultures for fermented foods of the future. *Int J Food Microbiol* 2002;**78**:119–31.
- Ben Said L, Gaudreau H, Dallaire L et al. Bioprotective culture: a new generation of food additives for the preservation of food quality and safety. *Ind Biotechnol* 2019;**15**:138–47. DOI: 10.1089/ind.2019.29175.lbs.
- Berg RD, Garlington AW. Translocation of certain indigenous bacteria from the gastrointestinal tract to the mesenteric lymph nodes and other organs in a gnotobiotic mouse model. *Infect Immun* 1979;**23**:403–11.
- Berg RD. Bacterial translocation from the gastrointestinal tract. *Trends Microbiol* 1995;**3**:149–54.
- Berg RD. Bacterial translocation from the intestines. *Jikken Dobutsu* 1985;**34**:1–16.
- Bernardeau M, Guguen M, Vernoux J. Beneficial lactobacilli in food and feed: long-term use, biodiversity and proposals for specific and realistic safety assessments. *FEMS Microbiol Rev* 2006;**30**:487–513.
- Borriello SP, Hammes WP, Holzapfel W et al. Safety of probiotics that contain lactobacilli or bifidobacteria. *Clin Infect Dis* 2003;**36**:775–80.
- Bosse Née Danz R, Müller A, Gibis M et al. Recent advances in cured raw ham manufacture. *Crit Rev Food Sci Nutr* 2018;**58**:610–30.
- Bourdichon F, Casaregola S, Farrokh C et al. Food fermentations: microorganisms with technological beneficial use. *Int J Food Microbiol* 2012;**154**:87–97. DOI: 10.1016/j.ijfoodmicro.2011.12.030. Epub 2011 Dec 31. Erratum in: *Int J Food Microbiol*. 2012 Jun 1;**156**(3):301. PMID: 22257932.
- Bover-Cid S, Holzapfel WF. Improved screening procedure for biogenic amine production by lactic acid bacteria. *Int J Food Microbiol* 1999;**53**:33–41.
- Bover-Cid S, Izquierdo-Pulido M, Vidal-Carou MC. Mixed starter cultures to control biogenic amine production in dry fermented sausages. *J Food Prot* 2000;**63**:1556–62.
- Callon C, Arliguie C, Montel MC. Control of Shigatoxin-producing *Escherichia coli* in cheese by dairy bacterial strains. *Food Microbiol* 2016;**53**:63–70.
- Callon C, Retureau E, Didienne R et al. Microbial biodiversity in cheese consortia and comparative *Listeria* growth on surfaces of uncooked pressed cheeses. *Int J Food Microbiol* 2014;**174**:98–109.
- Caplice E, Fitzgerald GF. Food fermentations: role of microorganisms in food production and preservation. *Int J Food Microbiol* 1999;**50**:131–49.
- Castellano P, Pérez Ibarreche M, Blanco Massani M et al. Strategies for pathogen biocontrol using lactic acid bacteria and their metabolites: a focus on meat ecosystems and industrial environments. *Microorganisms* 2017;**5**:38.
- Cebrián E, Rodríguez M, Peromingo B et al. Efficacy of the combined protective cultures of *Penicillium chrysogenum* and *Debaryomyces hansenii* for the control of ochratoxin A hazard in dry-cured ham. *Toxins* 2019;**11**:710.
- Chavan RS, Chavan SR. Sourdough technology—a traditional way for wholesome foods: a review. *Comp Rev Food Sci Food Saf* 2011;**10**:170–83. DOI: 10.1111/j.1541-4337.2011.00148.x.
- Chhetri V, Prakitchaiwattana C, Settachaimongkon S. A potential protective culture; halophilic *Bacillus* isolates with bacteriocin encoding gene against *Staphylococcus aureus* in salt added foods. *Food Control* 2019;**104**:292–9.
- Cleusix V, Lacroix C, Vollenweider S et al. Inhibitory activity spectrum of reuterin produced by *Lactobacillus reuteri* against intestinal bacteria. *BMC Microbiol* 2007;**7**:101.
- Cocolin L, Foschino R, Comi G et al. Description of the bacteriocins produced by two strains of *Enterococcus faecium* isolated from Italian goat milk. *Food Microbiol* 2007;**24**:752–8.

- Corsetti A, M. Gobbetti M, Rossi J et al. Antimould activity of sourdough lactic acid bacteria: identification of a mixture of organic acids produced by *Lactobacillus sanfrancisco* CB1. *Appl Microbiol Biotechnol* 1998;**50**:253–6.
- Crowley S, Mahony J, van Sinderen D. Current perspectives on antifungal lactic acid bacteria as natural bio-preservatives. *Trends Food Sci Technol* 2013;**33**:93–109.
- Cui Y, Zhang C, Wang Y et al. Class IIa bacteriocins: diversity and new developments. *Int J Mol Sci* 2012;**13**:16668–707.
- Dal Bello F, Clarke CI, Ryan LAM et al. Improvement of the quality and shelf life of wheat bread by fermentation with the antifungal strain *Lactobacillus plantarum* FST 1.7. *J Cereal Sci* 2007;**45**: 309–18.
- De Kwaadsteniet M, Ten Doeschate K, Dicks LM. Characterization of the structural gene encoding nisin F, a new lantibiotic produced by a *Lactococcus lactis* subsp. *lactis* isolate from freshwater catfish (*Clarias gariepinus*). *Appl Environ Microbiol* 2008;**74**:547–9.
- De Vos WM, Mulders JW, Siezen RJ et al. Properties of nisin Z and distribution of its gene, *nisZ*, in *Lactococcus lactis*. *Appl Environ Microbiol* 1993;**59**:213–8.
- De Vuyst L, Leroy F. Bacteriocins from lactic acid bacteria: production, purification, and food applications. *J Mol Microbiol Biotechnol* 2007;**13**:194–9.
- Delgado J, Núñez F, Asensio M.A et al. Quantitative proteomic profiling of ochratoxin A repression in *Penicillium nordicum* by protective cultures. *Int J Food Microbiol* 2019;**305**:108243.
- Dieuleveux V, Lemarinier S, Guéguen M. Antimicrobial spectrum and target site of d-3-phenyllactic acid. *Int J Food Microbiol* 1998;**40**:177–83.
- Dishisha T, Pereyra LP, Pyo SH et al. Flux analysis of the *Lactobacillus reuteri* propanediol-utilization pathway for production of 3-hydroxypropionaldehyde, 3-hydroxypropionic acid and 1,3-propanediol from glycerol. *Microb Cell Fact* 2014;**13**:76.
- Dong Q, Zhang W, Guo L et al. Influence of *Lactobacillus plantarum* individually and in combination with low O₂-MAP on the pathogenic potential of *Listeria monocytogenes* in cabbage. *Food Control* 2020;**107**:106765. DOI: 10.1016/j.foodcont.2019.106765.
- Doores S. Organic acids. In B. A. L. Davidson P.M. (Ed.), *Antimicrobials in Foods*. 3rd edn Marcel Dekker Inc, New York, NY, 2005, 95–136.
- Drosinos EH, Mataragas M, Metaxopoulos J. Biopreservation: a new direction towards food safety. In *New Developments in Food Policy, Control and Research*. ed. Riley, A.P. Nova Science Publishers. 2005.
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Ricci A, Allende AEFSa BIOHAZ Panel (EFSA Panel on Biological Hazards) et al. Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 7: suitability of taxonomic units notified to EFSA until September 2017. *EFSA Journal* 2018;**16**:5131, 43 pp.
- Elsser-Gravesen D, Elsser-Gravesen A. Biopreservatives. *Adv Biochem Eng Biotechnol* 2014;**143**:29–49.
- Enache-Angoulvant A, Hennequin C. Invasive *Saccharomyces* infection: a comprehensive review. *Clin Infect Dis* 2005;**41**:1559–68.
- Farnworth ER. Handbook of fermented functional foods. In: Mazza G (ed.). *Functional Foods and Nutraceuticals Series*, 2nd edn. Boca Raton: CRC Press Taylor & Francis Group. 2008.
- FEEDAP. Opinion of the scientific panel on additives and products or substances used in animal feed (FEEDAP) on the updating of the criteria used in the assessment of bacteria for resistance to antibiotics of human or veterinary importance. *EFSA J* 2005;**223**:1–12.
- FEEDAP. Technical guidance - Update of the criteria used in the assessment of bacterial resistance to antibiotics of human or veterinary importance. *EFSA J* 2008;**732**:1–15.
- Gaggia F, Di Gioia D, Baffoni L et al. The role of protective and probiotic cultures in food and feed and their impact on food safety. *Trends Food Sci Technol* 2011;**22**:S58–66.
- Garai-Ibabe G, Ibarburu I, Berregi I et al. Glycerol metabolism and bitterness producing lactic acid bacteria in cidermaking. *Int J Food Microbiol* 2008;**121**:253–61.
- Garde S, Avila M, Arias R et al. Outgrowth inhibition of *Clostridium beijerinckii* spores by a bacteriocin-producing lactic culture in ovine milk cheese. *Int J Food Microbiol* 2011;**150**: 59–65.
- Ge X, Teng K, Wang J et al. Identification of Key Residues in the NisK Sensor Region for Nisin Biosynthesis Regulation. *Front Microbiol* 2017;**8**:106.
- Giaouris E, Heir. E, Desvaux M et al. Intra- and inter-species interactions within biofilms of important foodborne bacterial pathogens. *Front Microbiol* 2015;**6**:841.
- Gross E, Morell JL. The structure of nisin. *J Am Chem Soc* 1971;**93**:4634–5.
- Gueimonde M, Ouwehand AC, Salminen S. Safety of Probiotics. *Scand J Nutr* 2005;**48**:42–8.
- Guimaraes A, Abrunhosa L, Pastrana L.M et al. Edible films and coatings as carriers of living microorganisms: a new strategy towards biopreservation and healthier foods. *Comp Rev Food Sci Food Saf* 2018;**17**:594–614.
- Habimana O, Meyrand M, Meylheuc T et al. Genetic features of resident biofilms determine attachment of *Listeria monocytogenes*. *Appl Environ Microbiol* 2009;**75**:7814–21.
- Hammes WP, Tichaczek PS. The potential of lactic acid bacteria for the production of safe and wholesome food (Das Potential der Milchsäurebakterien bei der Produktion von hygienisch sicheren und gesundheitsfördernden Lebensmitteln). *Zeitschrift für Lebensmittel-Untersuchung und -Forschung* 1994;**198**:193–201.
- Health Canada. Amendment (Schedule No. 948) to Division 28 of the Food and Drug Regulations, Sections B.28.001–003. Food and Drugs Regulations. 2003.
- Hibbing ME, Fuqua C, Parsek M.R et al. Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 2010;**8**:15–25.
- Honoré AH, Aunsbjerg SD, Ebrahimi P et al. Metabolic footprinting for investigation of antifungal properties of *Lactobacillus paracasei*. *Anal Bioanal Chem* 2016;**408**:83–96.
- Horowitz HW, Handwerker S, van Horn KG et al. *Leuconostoc*, an emerging vancomycin-resistant pathogen. *Lancet North Am Ed* 1987;**330**:1329–30.
- Husni RN, Gordon SM, Washington J.A et al. *Lactobacillus bacteremia* and endocarditis: review of 45 cases. *Clin Infect Dis* 1997;**25**:1048–55.
- Imran M, Bré JM, Guéguen M et al. Reduced growth of *Listeria monocytogenes* in two model cheese microcosms is not associated with individual microbial strains. *Food Microbiol* 2013;**33**:30–39.
- Jacques N, Casaregola S. Safety assessment of dairy microorganisms: the hemiascomycetous yeasts. *Int J Food Microbiol* 2008;**126**:321–6.
- Kandler O. Carbohydrate metabolism in lactic acid bacteria. *Antonie Van Leeuwenhoek* 1983;**49**:209–24.
- Kawtharani H, Snini SP, Heang S et al. Phenyllactic acid produced by *Geotrichum candidum* reduces *Fusarium sporotrichioides* and

- F. langsethiae* growth and T-2 toxin concentration. *Toxins* 2020;12:209.
- Kim DH, Jeong D, Kim H et al. Antimicrobial activity of Kefir against various food pathogens and spoilage bacteria. *Kor J Food Sci Anim Resour* 2016;36:787–90.
- Kjos M, Oppgaard C, Diep DB et al. Sensitivity to the two-peptide bacteriocin lactococcin G is dependent on UppP, an enzyme involved in cell-wall synthesis. *Mol Microbiol* 2014;92:1177–87.
- Kondrotiene K, Kasnauskaitė N, Serniene L et al. Characterization and application of newly isolated nisin producing *Lactococcus lactis* strains for control of *Listeria monocytogenes* growth in fresh cheese. *LWT Food Sci Technol* 2018;87:507–14.
- Kuipers OP, Beerthuyzen MM, Siezen RJ et al. Characterization of the nisin gene cluster nisABTCIPR of *Lactococcus lactis*. Requirement of expression of the nisA and nisI genes for development of immunity. *Eur J Biochem* 1993;216:281–91.
- Lacroix N, St Gelais D, Champagne CP et al. Characterization of aromatic properties of old-style cheese starters. *J Dairy Sci* 2010;93:3427–41.
- Laranjo M, Potes ME, Elias M. Role of starter cultures on the safety of fermented meat products. *Front Microbiol* 2019;10:853.
- Laulund S, Wind A, Derkx PMF et al. Regulatory and safety requirements for food cultures. *Microorganisms* 2017;5:28.
- Lavermicocca P, Valerio F, Evidente A et al. Purification and characterization of novel antifungal compounds from the sourdough *Lactobacillus plantarum* strain 21B. *Appl Environ Microbiol* 2000;66:4084–90.
- Lavermicocca P, Valerio F, Visconti A. Antifungal activity of phenyllactic acid against molds isolated from bakery products. *Appl Environ Microbiol* 2003;69:634–40.
- Lee JY, Shim JM, Yao Z et al. Antimicrobial activity of *Bacillus amyloliquefaciens* EMD17 isolated from Cheonggukjang and potential use as a starter for fermented soy foods. *Food Sci Biotechnol* 2016;25:525–32.
- Leuschner R-GK, Robinson T, Hugas M et al. Qualified presumption of safety (QPS): a generic risk assessment approach for biological agents notified to the European Food Safety Authority (EFSA). *Trends Food Sci Technol* 2010;21:425–35.
- Leyva Salas M, Mounier J, Valence F et al. Antifungal microbial agents for food biopreservation—a review. *Microorganisms* 2017;5:37.
- Leyva Salas M, Thierry A, Lemaître M et al. Antifungal activity of lactic acid bacteria combinations in dairy mimicking models and their potential as bioprotective cultures in pilot scale applications. *Front Microbiol* 2018;9:1787.
- Li L, Shin SY, Lee KW et al. Production of natural antimicrobial compound D-phenyllactic acid using *Leuconostoc mesenteroides* ATCC 8293 whole cells involving highly active D-lactate dehydrogenase. *Lett Appl Microbiol* 2014;59:404–11.
- Li X, Jiang B, Pan B et al. Purification and partial characterization of *Lactobacillus* species SK007 lactate dehydrogenase (LDH) catalyzing phenylpyruvic acid (PPA) conversion into phenyllactic acid (PLA). *J Agric Food Chem* 2008;56:2392–9.
- Li X, Jiang B, Pan B. Biotransformation of phenylpyruvic acid to phenyllactic acid by growing and resting cells of a *Lactobacillus* sp. *Biotechnol Lett* 2007;29:593–7.
- Lianou A, Samelis J. Addition to thermized milk of *Lactococcus lactis* subsp. *cremoris* M104, a wild, novel nisin a-producing strain, replaces the natural antilisterial activity of the autochthonous raw milk microbiota reduced by thermization. *J Food Prot* 2014;77:1289–97.
- Lin XB, Gänzle MG. Effect of lineage-specific metabolic traits of *Lactobacillus reuteri* on sourdough microbial ecology. *Appl Environ Microbiol* 2014;80:5782–9.
- Liong MT. Safety of probiotics: translocation and infection. *Nutr Rev* 2008;66:192–202.
- Lubelski J, Rink R, Khusainov R et al. Biosynthesis, immunity, regulation, mode of action and engineering of the model lantibiotic nisin. *Cell Mol Life Sci* 2008;65:455–76.
- Magnusson J, Strom K, Roos S et al. Broad and complex antifungal activity among environmental isolates of lactic acid bacteria. *FEMS Microbiol Lett* 2003;219:129–35.
- Makarova K, Slesarev A, Wolf Y et al. Comparative genomics of the lactic acid bacteria. *Proc Natl Acad Sci* 2006;103:15611–6.
- Maligoy M, Mercade M, Coccagn-Bousquet M et al. *Appl Environ Microbiol* 2008;74:485–94.
- Mani-López E, García HS, López-Malo A. Organic acids as antimicrobials to control *Salmonella* in meat and poultry products. *Food Res Int* 2012;45:713–21.
- Mannazzu I, Domizio P, Carboni G et al. Yeast killer toxins: from ecological significance to application. *Crit Rev Biotechnol* 2019. 10.1080/07388551.2019.1601679.
- Marilley L, Casey MG. Flavors of cheese products: metabolic pathways, analytical tools and identification of producing strains. *Int J Food Microbiol* 2004;90:139–59.
- Martín R, Olivares M, Marín ML et al. Characterization of a reuterin-producing *Lactobacillus coryniformis* strain isolated from a goat's milk cheese. *Int J Food Microbiol* 2005;104:267–77.
- Martinez RCR, Staliano CD, Vieira ADS et al. Bacteriocin production and inhibition of *Listeria monocytogenes* by *Lactobacillus sakei* subsp *sakei* 2a in a potentially synbiotic cheese spread. *Food Microbiol* 2015;48:143–52.
- McAuliffe O, Ross RP, Hill C. Lantibiotics: structure, biosynthesis and mode of action. *FEMS Microbiol Rev* 2001;25:285–308.
- Medina-Córdova N, Rosales-Mendoza S, Hernández-Montiel LG et al. The potential use of *Debaryomyces hansenii* for the biological control of pathogenic fungi in food. *Biol Control* 2018;121:216–22.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infect Dis* 2011;11:142–51.
- Mirkovic N, Radulovic Z, Uzelac G et al. Isolation and characterisation of bacteriocin and aggregation-promoting factor production in *Lactococcus lactis* ssp. *lactis* BGBM50 strain. *Food Technol Biotechnol* 2015;53:237–42.
- Mofredj A, Bahloul H, Chanut C. *Lactococcus lactis*: an opportunistic bacterium? *Médecine et Maladies Infectieuses* 2007;37:200–7.
- Moll G, Hauge HH, Nissen-Meyer J et al. Mechanistic properties of the two component bacteriocin lactococcin G. *J Bacteriol* 1998;180:96–99.
- Moll G, Ubbink-Kok T, Hildeng-Hauge H et al. Lactococcin G is a potassium ion-conducting, two component bacteriocin. *J Bacteriol* 1996;178:600–5.
- Monnet C, Back A, Irlinger F. Growth of aerobic ripening bacteria at the cheese surface is limited by the availability of iron. *Appl Environ Microbiol* 2012;78:3185–92.
- Morata A, Escott C, Banuelos A et al. Contribution of Non-Saccharomyces yeasts to wine freshness. A review. *Biomolecules* 2020;10:34.
- Morita H, Toh H, Fukuda S et al. Comparative genome analysis of *Lactobacillus reuteri* and *Lactobacillus fermentum* reveal a genomic island for reuterin and cobalamin production. *DNA Res* 2008;15:151–61.
- Motarjemi Y, Nout MJR, Joint FAO/WHO Workshop on Assessment of Fermentation as a Household Technology for

- Improving Food Safety. Food fermentation: a safety and nutritional assessment. *Bull World Health Organ* 1996;**74**: 553–9.
- Mu W, Yang Y, Jia J et al. Production of 4-hydroxyphenyllactic acid by *Lactobacillus* sp. SK007 fermentation. *J Biosci Bioeng* 2010;**109**:369–71.
- Muhialdin BJ, Algboory HL, Kadum H et al. Antifungal activity determination for the peptides generated by *Lactobacillus plantarum* TE10 against *Aspergillus flavus* in maize seeds. *Food Control* 2020;**109**:106898.
- Nawaz M, Wang J, Zhou A et al. Characterization and transfer of antibiotic resistance in lactic acid bacteria from fermented food products. *Curr Microbiol* 2011;**62**:1081–9.
- Nazareth TM, Luz C, Torrijos R et al. Potential application of lactic acid bacteria to reduce Aflatoxin B1 and Fumonisin B1 occurrence on corn kernels and corn ears. *Toxins* 2020;**12**:21.
- Ndagano D, Lamoureux T, Dortu C et al. Antifungal activity of 2 lactic acid bacteria of the *Weissella* genus isolated from food. *J Food Sci* 2011;**76**:305–11.
- Nes IF, Diep DB, Håvarstein LS et al. Biosynthesis of bacteriocins in lactic acid bacteria. *Antonie Van Leeuwenhoek* 1996;**70**: 113–28.
- Nikodinoska I, Baffoni L, Di Gioia D et al. Protective cultures against foodborne pathogens in a nitrite reduced fermented meat product. *LWT* 2019;**101**:293–9.
- Niku-Paavola ML, Laitila A, Mattila-Sandholm T et al. New types of antimicrobial compounds produced by *Lactobacillus plantarum*. *J Appl Microbiol* 1999;**86**:29–35.
- Nissen-Meyer J, Holo H, Håvarstein LS et al. A novel lactococcal bacteriocin whose activity depends on the complementary action of two peptides. *J Bacteriol* 1992;**174**:5686–92.
- O'Sullivan JN, O'Connor PM, Rea MC et al. Nisin J, a novel natural Nisin variant, is produced by *Staphylococcus capitis* sourced from the human skin microbiota. *J Bacteriol* 2020;**202**: e00639–19.
- Oliveira M, Ferreira V, Magalhães R et al. Biocontrol strategies for Mediterranean-style fermented sausages. *Food Res Int* 2018;**103**:438–49.
- Oppgaard C, Emanuelsen L, Thorbek L et al. The lactococcal G immunity protein recognizes specific regions in both peptides constituting the two-peptide bacteriocin lactococcin G. *Appl Environ Microbiol* 2010;**76**:1267–73.
- Poutanen K, Flander L, Katina K Sourdough and cereal fermentation in a nutritional perspective. *Food Microbiol* 2009;**26**:693–9.
- Prema P, Smila D, Palavesam A et al. Production and characterization of an antifungal compound (3-phenyllactic acid) produced by *Lactobacillus plantarum* strain. *Food Bioprocess Technol* 2010;**3**:379–86.
- Ramos B, Brandão TRS, Teixeira P et al. Biopreservation approaches to reduce *Listeria monocytogenes* in fresh vegetables. *Food Microbiol* 2020;**85**:103282.
- Raspor P, Goranovič D. Biotechnological applications of acetic acid bacteria. *Crit Rev Biotechnol* 2008;**28**:101–24.
- Rodríguez JM, Martínez MI, Kok J. Pediocin PA-1, a wide-spectrum bacteriocin from lactic acid bacteria. *Crit Rev Food Sci Nutr* 2002;**42**:91–121.
- Rodríguez N, Salgado JM, Cortés S et al. Antimicrobial activity of d-3-phenyllactic acid produced by fed-batch process against *Salmonella enterica*. *Food Control* 2012;**25**:274–84.
- Rogne P, Fimland G, Nissen-Meyer J et al. Three-dimensional structure of the two peptides that constitute the two-peptide bacteriocin lactococcin G. *Biochim Biophys Acta* 2008;**1784**:543–54.
- Ross RP, Morgan S, Hill C. Preservation and fermentation: past, present and future. *Int J Food Microbiol* 2002;**79**:3–16.
- Roth E, Schwenninger SM, Eugster-Meier E et al. Facultative anaerobic halophilic and alkaliphilic bacteria isolated from a natural smear ecosystem inhibit *Listeria* growth in early ripening stages. *Int J Food Microbiol* 2011;**147**:26–32.
- Rouse S, van Sinderen D. Bioprotective potential of lactic acid bacteria in malting and brewing. *J Food Prot* 2008;**71**:1724–33.
- Russo P, Fares C, Longo A et al. *Lactobacillus plantarum* with broad antifungal activity as a protective starter culture for bread production. *Foods* 2017;**6**:1101–9.
- Ryan LAM, Zannini E, Bello DF et al. *Lactobacillus amylovorus* DSM 19280 as a novel food-grade antifungal agent for bakery products. *Int J Food Microbiol* 2011;**146**:276–83.
- Saarela M, Maukonen J, von Wright A et al. Tetracycline susceptibility of the ingested *Lactobacillus acidophilus* LaCH-5 and *Bifidobacterium animalis* subsp. *lactis* Bb-12 strains during antibiotic/probiotic intervention. *Int J Antimicrob Agents* 2007;**29**:271–80.
- Salas ML, Mounier J, Valence F et al. Antifungal microbial agents for food biopreservation. A review. *Microorganisms* 2017;**5**:37.
- Samelis J, Kakouri A. Hurdle factors minimizing growth of *Listeria monocytogenes* while counteracting in situ antilisterial effects of a novel nisin A-producing *Lactococcus lactis* subsp. *cremoris* costarter in thermized cheese milks. *AIMS Microbiol* 2018;**4**:19–41.
- Santos F, Vera JL, van der Heijden R et al. The complete coenzyme B12 biosynthesis gene cluster of *Lactobacillus reuteri* CRL1098. *Microbiology* 2008;**154**:81–93.
- Saravanakumar D, Ciavarella A, Spadaro D et al. *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and *Penicillium expansum* in apples through iron depletion. *Postharvest Biol Technol* 2008;**49**: 121–8.
- Sauvageot N, Gouffi K, Laplace JM et al. Glycerol metabolism in *Lactobacillus collinoides*: production of 3-hydroxypropionaldehyde, a precursor of acrolein. *Int J Food Microbiol* 2000;**55**:167–70.
- Sauvageot N, Muller C, Hartke A et al. Characterisation of the diol dehydratase pdu operon of *Lactobacillus collinoides*. *FEMS Microbiol Lett* 2002;**209**:69–74.
- Scatassa ML, Gaglio R, Cardamone C et al. Anti-listeria activity of lactic acid bacteria in two traditional Sicilian cheeses. *Ital J Food Saf* 2017;**6**:6191.
- Schaefer L, Auchtung TA, Hermans KE et al. The antimicrobial compound reuterin (3-hydroxypropionaldehyde) induces oxidative stress via interaction with thiol groups. *Microbiol* 2010;**156**:1589–99.
- Schütz H, Radler F. Anaerobic reduction of glycerol to propanediol-1.3 by *Lactobacillus brevis* and *Lactobacillus buchneri*. *Syst Appl Microbiol* 1984;**5**:169–78.
- Semedo T, Almeida SM, Martins P et al. Comparative study using type strains and clinical and food isolates to examine hemolytic activity and occurrence of the *cyl* operon in *Enterococci*. *J Clin Microbiol* 2003a;**41**:2569–76.
- Semedo T, Santos MA, Lopes MF et al. Virulence factors in food, clinical and reference *Enterococci*: a common trait in the genus? *Syst Appl Microbiol* 2003b;**26**:13–22.
- Settanni L, Corsetti A. Application of bacteriocins in vegetable food biopreservation. *Int J Food Microbiol* 2008;**121**:123–38.
- Settanni L, Franciosi E, Cavazza A et al. Extension of Tosèla cheese shelf-life using non-starter lactic acid bacteria. *Food Microbiol* 2011;**28**:883–90.

- Sicard D, Legras JL. Bread, beer and wine: yeast domestication in the *Saccharomyces sensu strict* complex. *C R Biol* 2011;**334**: 229–36.
- Siedler S, Balti R, Neves AR. Bioprotective mechanisms of lactic acid bacteria against fungal spoilage of food. *Curr Opin Biotechnol* 2019;**56**:138–46.
- Siedler S, Rau MH, Bidstrup S et al. Competitive exclusion is a major bioprotective mechanism of *Lactobacilli* against fungal spoilage in fermented milk products. *Appl Environ Microbiol* 2020;**86**:e02312–19.
- Sieuwerds S, de Bok FAM, Hugenholtz J et al. Unraveling microbial interactions in food fermentations: from classical to genomics approaches. *Appl Environ Microbiol* 2008;**74**: 4997–5007.
- Silva CCG, Silva SPM, Ribeiro SC. Application of bacteriocins and protective cultures in dairy food preservation. *Front Microbiol* 2018;**9**:594. <https://www.frontiersin.org/article/10.3389/fmicb.2018.00594>.
- Sindi A, Badsha MB, Nielsen B et al. Antimicrobial activity of six international Artisanal Kefirs against *Bacillus cereus*, *Listeria monocytogenes*, *Salmonella enterica* serovar Enteritidis, and *Staphylococcus aureus*. *Microorganisms* 2020;**8**:849.
- Sipiczki M. *Metschnikowia* strains isolated from botrytized grapes antagonize fungal and bacterial growth by iron depletion. *Appl Environ Microbiol* 2006;**72**:6716–24.
- Smit G, Smit BA, Engels WJ. Flavor formation by lactic acid bacteria and biochemical flavor profiling of cheese products. *FEMS Microbiol Rev* 2005;**29**:591–610.
- Spano G, Russo P, Lonvaud-Funel A et al. Biogenic amines in fermented foods. *Eur J Clin Nutr* 2010;**64**:S95–100.
- Sriramulu DD, Liang M, Hernandez-Romero D et al. *Lactobacillus reuteri* DSM 20016 produces cobalamin-dependent diol dehydratase in metabolosomes and metabolizes 1,2-propanediol by disproportionation. *J Bacteriol* 2008;**190**:4559–67.
- Starmer WT, Ganter PF, Aberdeen V et al. The ecological role of killer yeasts in natural communities of yeasts. *Can J Microbiol* 1987;**33**:783–96.
- Straub BW, Kicherer M, Schilcher SM et al. The formation of biogenic amines by fermentation organisms (Die Bildung von biogenen Aminen durch Fermentationsorganismen). *Zeitschrift für Lebensmittel-Untersuchung und Forschung* 1995;**201**:79–82.
- Ström K, Sjögren J, Broberg A et al. *Lactobacillus plantarum* MiLAB 393 produces the antifungal cyclic dipeptides cyclo(l-Phe-l-Pro) and cyclo(l-Phe-trans-4-OH-l-Pro) and phenyl lactic acid. *Appl Environ Microbiol* 2002;**68**:4322–7.
- Sumner SS, Speckhard MW, Somers EB et al. Isolation of histamine-producing *Lactobacillus buchneri* from Swiss cheese implicated in a food poisoning outbreak. *Appl Environ Microbiol* 1985;**50**:1094–6.
- Sun D, Li H, Song D et al. Genome, transcriptome and fermentation analyses of *Lactobacillus plantarum* LY-78 provide new insights into the mechanism of phenyllactate biosynthesis in lactic acid bacteria. *Biochem Biophys Res Commun* 2019;**519**:351–7.
- Talarico TL, Dobrogosz WJ. Purification and characterization of glycerol dehydratase from *Lactobacillus reuteri*. *Appl Environ Microbiol* 1990;**56**:1195–7.
- Tarazanova M, Beerthuyzen M, Siezen R et al. Plasmid complement of *Lactococcus lactis* NCDO712 reveals a novel pilus gene cluster. *PLoS ONE* 2016;**11**:e0167970.
- Valerio F, Di Biase M, Lattanzio VMT et al. Improvement of the antifungal activity of lactic acid bacteria by addition to the growth medium of phenylpyruvic acid, a precursor of phenyllactic acid. *Int J Food Microbiol* 2016;**222**: 1–7.
- Valerio F, Lavermicocca P, Pascale M et al. Production of phenyllactic acid by lactic acid bacteria: an approach to the selection of strains contributing to food quality and preservation. *FEMS Microbiol Lett* 2004;**233**:289–95.
- Van Boekel M, Fogliano V, Pellegrini N et al. A review on the beneficial aspects of food processing. *Mol Nutr Food Res* 2010;**54**:1215–47.
- Van Gijtenbeek LA, Eckhardt TH, Herrera-Domínguez L et al. Gene-trait matching and prevalence of Nisin tolerance systems in *Lactococcus lactis*. *Front Bioeng Biotechnol* 2021;**9**: 622835.
- Vermeulen N, Ganzle MG, Vogel RF. Influence of peptide supply and cosubstrates on phenylalanine metabolism of *Lactobacillus sanfranciscensis* DSM20451T and *Lactobacillus plantarum* TMW1.468. *J Agric Food Chem* 2006;**54**: 3832–9.
- Vollenweider S, Grassi G, König I et al. Purification and structural characterization of 3-hydroxypropionaldehyde and its derivatives. *J Agric Food Chem* 2003;**51**: 3287–93.
- Vorobjeva L, Khodjaev EY, Vorobjeva NV. Propionic acid bacteria as probiotic. *Microb Ecol Health Dis* 2008;**20**:109–12.
- Wels M, Siezen R, van Hijum S et al. Comparative genome analysis of *Lactococcus lactis* indicates niche adaptation and resolves genotype/phenotype disparity. *Front Microbiol* 2019;**10**:4.
- Wilson PDG, Wilson DR, Waspe CR. Weak acids: dissociation in complex buffering systems and partitioning into oils. *J Sci Food Agric* 2000;**80**:471–6.
- Wirawan RE, Klesse NA, Jack RW et al. Molecular and genetic characterization of a novel nisin variant produced by *Streptococcus uberis*. *Appl Environ Microbiol* 2006;**72**: 1148–56.
- Wu W, Deng G, Liu C et al. Optimization and multiomic basis of phenyllactic acid overproduction by *Lactobacillus plantarum*. *J Agric Food Chem* 2020;**68**:1741–9.
- Yamada Y, Yukphan P. Genera and species in acetic acid bacteria – a review. *Int J Food Microbiol* 2008;**125**:15–24.
- Yvon M, Thirouin S, Rijnen L et al. An aminotransferase from *Lactococcus lactis* initiates conversion of amino acids to cheese flavor compounds. *Appl Environ Microbiol* 1997;**63**: 414–9.
- Zendo T, Fukao M, Ueda K et al. Identification of the lantibiotic nisin Q, a new natural nisin variant produced by *Lactococcus lactis* 61-14 isolated from a river in Japan. *Biosci Biotechnol Biochem* 2003;**67**:1616–9.
- Zhang P, Ganzle M, Yang X. Complementary antibacterial effects of bacteriocins and organic acids as revealed by comparative analysis of *Carnobacterium* spp. from meat. *Appl Environ Microbiol* 2019;**85**:e01227–01219.