

OCADIA

Hybrid Ryegrass (4n)

Lolium x hybridum Hausskn.

Things to know

The variety Ocadia belongs to the intermediate type of hybrid ryegrass and it forms considerably less stems in the estival growth than the Italian type like the variety Rusa. Thereby, Ocadia has a higher digestibility. Ocadia provides high yields and shows a very good endurance. Its resistances against rust fungi and bacterial wilt are excellent.

Descent

Base material

Selection within breeding material from Reckenholz, emerged from a colchicine treatment.

M0 seed

Row trial 2001 (LH0105) with seed harvest on 4 clone progenies from a polycross with 6 clones.

Literature

Frick R.,Mosimann E.,Suter D.,Hirschi H.-U., 2010. Bastard-Raigras und Wiesen-Fuchsschwanz: Sortenversuche 2007 bis 2009. Agrarforschung Schweiz 1(9), 334-339

National listing

Situation in Switzerland

On the Swiss List of Recommended Varieties of Forage Plants since 2010

Situation abroad

FR (Representative: Semences Vertes)

Agronomic caracteristics

Results of the official Swiss variety trials 2007-2009

(Frick et al. 2010)

(FIICK & al. 2010)	OCADIA	Mean
Yield	3.5	4.3
General impression	4.4	4.7
Juvenile growth	3.9	4.0
Competing ability	4.7	4.8
Persistence	4.7	5.1
Resistance to winter conditions	5.4	5.1
Resistance to leafspots an rust	3.1	3.2
Resistance to bacterial wilt	1.5	2.2
Digestibility of the organic matter	5.0	5.1
Index (weighted average of all notes)	4.0	4.3

Scoring scale 1 = very good; 5 = medium; 9 = very poor Yield Mean of 4 experimental sites over 2 years Mean value of standard varieties

Description according to UPOV gidelines

DUS test conducted at Crossnacreevy, DARD (UK), 2006-2008

UPOV No Characteristics State of expression 1 Ploidy tetraploid 2 Plant: growth habit semi-prostrate 3 Tendency to form inflorescences (without vernalization) 8 Time of inflorescence emergence (after vernalization) 10 Flag leaf: length long 11 Flag leaf: width medium 12 Stem: length of longest stem including inflorescence	
2 Plant: growth habit semi-prostrate 3 Tendency to form inflorescences (without vernalization) 8 Time of inflorescence emergence (after vernalization) 10 Flag leaf: length long 11 Flag leaf: width medium 12 Stem: length of longest stem including medium	Note
Tendency to form inflorescences (without vernalization) Time of inflorescence emergence (after vernalization) Flag leaf: length long Flag leaf: width medium Stem: length of longest stem including medium	4
vernalization) Time of inflorescence emergence (after vernalization) Flag leaf: length long Flag leaf: width medium Stem: length of longest stem including medium	7
vernalization) 10 Flag leaf: length long 11 Flag leaf: width medium 12 Stem: length of longest stem including medium	1
11 Flag leaf: width medium 12 Stem: length of longest stem including medium	3
12 Stem: length of longest stem including medium	7
	5
irinorescence	5

Version: 06.06.2012

Publisher: Agroscope Reckenholz-Tänikon Research Station ART,

Reckenholzstrasse 191, 8046 Zürich

In Collaboration with: Delley Seeds and Plants Ltd (DSP), 1567 Delley Editorial Team: Beat Boller, Peter Tanner, Lucienne Adam, ART

Copyright: © 2012, Art





Schweizerische Eidgenossenschaft Confédération suisse Confederazione Svizzera Confederaziun svizra

Swiss Confederation

Federal Department of Economic Affairs FDEA Agroscope Reckenholz-Tänikon Research Station ART