



Biocontrol yeasts: mechanisms and applications

Florian M. Freimoser¹ · Maria Paula Rueda-Mejia¹ · Bruno Tilocca^{2,4} · Quirico Migheli^{2,3}

Received: 17 July 2019 / Accepted: 17 September 2019
© The Author(s) 2019

Abstract

Yeasts occur in all environments and have been described as potent antagonists of various plant pathogens. Due to their antagonistic ability, undemanding cultivation requirements, and limited biosafety concerns, many of these unicellular fungi have been considered for biocontrol applications. Here, we review the fundamental research on the mechanisms (e.g., competition, enzyme secretion, toxin production, volatiles, mycoparasitism, induction of resistance) by which biocontrol yeasts exert their activity as plant protection agents. In a second part, we focus on five yeast species (*Candida oleophila*, *Aureobasidium pullulans*, *Metschnikowia fructicola*, *Cryptococcus albidus*, *Saccharomyces cerevisiae*) that are or have been registered for the application as biocontrol products. These examples demonstrate the potential of yeasts for commercial biocontrol usage, but this review also highlights the scarcity of fundamental studies on yeast biocontrol mechanisms and of registered yeast-based biocontrol products. Yeast biocontrol mechanisms thus represent a largely unexplored field of research and plentiful opportunities for the development of commercial, yeast-based applications for plant protection exist.

Keywords Biological control · Microbial antagonism · Plant protection · Plant pathogens · Competition · Enzyme secretion · Toxin production · Volatiles · Mycoparasitism · Resistance induction

Introduction

Despite their relevance as model eukaryotes for biotechnological applications and in medical mycology, the potential use of antagonistic yeasts as biocontrol agents is still underexploited. Only a handful of yeast-based plant protection products has reached the market and even in fundamental research, antifungal yeasts have been neglected and poorly investigated with state-of-the-art technology and

at the molecular level. Nevertheless, yeasts combine strong antifungal activities with advantageous properties for an application (e.g., strong antagonistic activity, culturability, formulatability, applicability, stress resistance) and are thereby promising for the development of biological plant protection agents. Furthermore, the close relatedness with model yeasts, particularly *Saccharomyces cerevisiae*, enables taking advantage of the molecular tools and plethora of data developed for these organisms for application-oriented and basic studies on biocontrol yeasts.

Biocontrol is mostly looked at and studied in a species/isolate-centric manner: different species/isolates are tested against the target plant pathogen and the most active organism is studied with respect to its potential for biocontrol applications. However, for a successful application and improvement of biocontrol organisms, we first have to understand the biocontrol mechanisms involved and then confirm their expression under field conditions (Droby and Chalutz 1994; Spadaro and Droby 2016; Wisniewski et al. 2007). Here, different yeast biocontrol mechanisms are highlighted and a comprehensive overview on published work on antagonistic mechanisms of biocontrol yeasts is provided in Supplementary Table 1.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11274-019-2728-4>) contains supplementary material, which is available to authorized users.

✉ Florian M. Freimoser
florian.freimoser@agroscope.admin.ch

- ¹ Agroscope, Research Division Plant Protection, Müller-Thurgau-Strasse 29, 8820 Wädenswil, Switzerland
- ² Dipartimento di Agraria, Università degli Studi di Sassari, Viale Italia 39, 07100 Sassari, Italy
- ³ Istituto Nazionale di Biostrutture e Biosistemi and NRD - Nucleo di Ricerca sulla Desertificazione, Università degli Studi di Sassari, Viale Italia 39, 07100 Sassari, Italy
- ⁴ Department of Health Sciences, University “Magna Græcia” of Catanzaro, Viale Europa, 88100 Catanzaro, Italy

Advantageous yeast properties for potential biocontrol applications

Any organism to be used as the active ingredient in a biocontrol product must be effective against its target disease, but secondary properties such as biosafety and registration issues, production requirements and conditions, formulation options, and the required application equipment are just as or even more important. Although the lack of invasive, filamentous growth of most yeasts may seem a disadvantage, the yeast-like morphology is the reason for widely culturability in fermentors, advantageous formulation characteristics and ample application options. As for bacteria, the single-celled morphology of yeasts also favours adhesion and biofilm formation, which directly influences environmental persistence, competitiveness and thereby improved biocontrol activity (Fanning and Mitchell 2012; Pandin et al. 2017; Rossouw et al. 2018; Verstrepen and Klis 2006).

While many *S. cerevisiae* strains contain the high-copy 2 µm plasmid (i.e., 653 of 1011 sequenced *S. cerevisiae* strains) (Peter et al. 2018), most non-conventional yeasts lack plasmids (but can be engineered to maintain foreign, extra-chromosomal DNA by designing a plasmid vector containing intrinsic autonomously replicating and centromere sequences) (Cao et al. 2017). Yeasts thus share growth characteristics and biocontrol activities with bacteria without the risk of taking up or passing on plasmid-based antibiotic resistance, pathogenicity factors or toxin biosynthesis genes. In addition, horizontal gene transfer, albeit occurring more frequently in fungi than thought earlier, is significantly less frequent in yeasts, as compared to their prokaryotic counterparts, due to their more complex genome organisation (Fitzpatrick 2012; Moriguchi et al. 2013; Richards et al. 2011).

Yeasts have been used for food and beverage production for thousands of years, they are consumed directly as food supplements, and are widely employed in the food industry (Bekatorou et al. 2006; Querol and Fleet 2006). In many cases, these “food industry yeasts” belong to the same genus or even species as those intended for biocontrol (e.g., *S. cerevisiae*, *Candida sake*, *Metschnikowia pulcherrima*). This may be the reason why yeasts *sensu lato* are generally regarded as safe and therefore applying yeasts in crops and on food products elicits less concern than applications of bacteria or filamentous fungi (European Food Safety Authority 2005). Nonetheless, some yeasts, such as certain *Candida* or *Cryptococcus* species, are important fungal human pathogens (Butler et al. 2009; Miceli et al. 2011; Opulente et al. 2019). Properties such as dimorphism (e.g., the switch to an invasive growth form), growth at high temperatures (e.g., at or

above 37 °C) and resistance to fungicides are of particular concern and should be studied and assessed in detail before considering new isolates for biocontrol applications (Gauthier 2015, 2017; Robert et al. 2015).

Mechanisms underlying the biocontrol activity of yeasts

Understanding the mechanisms conferring biocontrol activity is the foundation for the informed and successful development and application of yeasts as plant protection agents (Droby and Chalutz 1994; Spadaro and Droby 2016; Wisniewski et al. 2007). For the biocontrol yeasts so far studied in detail, multiple mechanisms such as competition for nutrients and space, secretion of enzymes, toxin production, release of volatile organic compounds (VOCs), myco-parasitism and induction of resistance in plants are likely to be involved in the antagonistic function (Fig. 1) (Droby et al. 2009; Punja and Utkhede 2003; Wisniewski and Droby 2012). In most cases, the mechanisms outlined and discussed below have not been fully proven by molecular analyses (e.g., by gene deletion and complementation, heterologous expression), but rather proposed based on analogies with other biological systems. However, the increasing number of annotated yeast genomes and the availability of different transformation techniques should make it possible to decipher different mechanisms and to unambiguously confirm biocontrol mechanisms in future work.




Fig. 1 Multiple mechanisms are involved in preventing plant diseases and conferring biocontrol activity to yeasts. The mechanisms studied and highlighted here are competition for nutrients and space, secretion of toxins, enzymes and volatile organic compounds, direct parasitisation (fungivory) and indirect mechanisms (i.e., induction of resistance)

Competition for nutrients and space

All microorganisms compete with each other and their hosts for nutrients and space: this struggle is considered as the primary mode of action of biocontrol yeasts (Schaible and Kaufmann 2005; Spadaro and Droby 2016; Wisniewski et al. 2007). Competition is difficult to study mechanistically: it is likely more important in natural environments, where resources are limited and competitors plentiful. In community ecology, niche and nutrient competition have been intensely studied as determinants of species diversity. In nectar yeasts (e.g., *Metschnikowia reukaufii*), which are closely related to some biocontrol species, efficient resource depletion, due to the duplication of nitrogen transporter and metabolism genes, causes priority effects (i.e., order of species arrival determines community composition) and thus acts as a driver of competitiveness among different species (Dhama et al. 2016). With respect to competition for space, *in vitro* experiments performed on solid media seem to ascribe a minor role to a limitation in space. Although most yeasts grow well on agar plates, large differences in their antifungal activities were observed (Hilber-Bodmer et al. 2017). In addition, species-specific inhibition does not seem to occur, and a particular yeast is either strongly or weakly antagonistic against most fungi (Hilber-Bodmer et al. 2017). However, growing under field conditions activates diverse survival mechanisms and the competition for the physical niche might gain importance in such circumstances.

Most organisms and cells, from humans to bacteria, synthesise iron binding molecules to deprive competing organisms, pathogens or intracellular parasites of this essential element (Barber and Elde 2015; Johnson 2008). Also for biocontrol yeasts, iron is one of the most sought after nutrients and the competition for iron is recognised as an important mode of action (Spadaro and Droby 2016). In *Aureobasidium pullulans*, a siderophore identified as fusarinine C (fusigen) was identified and shown to exhibit antibacterial activity (Wang et al. 2009a, b). The peculiar red color of *M. pulcherrima* colonies is due to the formation of a cyclic dipeptide, pulcherriminic acid, that complexes iron (Gore-Lloyd et al. 2019). Pigmentless *M. pulcherrima* mutants were shown to exhibit reduced antifungal activity and iron deprivation of the fungal pathogen was suggested as one of several mechanisms by which this yeast antagonises plant pathogenic fungi (Gore-Lloyd et al. 2019; Sipiczki 2006). However, mutants lacking the ability to synthesise pulcherriminic acid still inhibited filamentous fungi strongly, suggesting that the antifungal activity was not only due to iron deprivation (Gore-Lloyd et al. 2019). The exact contribution of iron chelators to yeast biocontrol activity thus remains to be elucidated in detail.

Recently, it was shown that *Saccharomycopsis schoenii* lacks several components of the sulfur assimilation pathway

and thus likely acquires methionine from its prey (Junker et al. 2019). Among yeasts, the inability to take up sulfur is specific to *Saccharomycopsis*, but some plant pathogenic fungi and *Trichoderma* species show a similar phenomenon, which may indicate that methionine is an important target for such organisms and highly competed over (Junker et al. 2019). Pioneering experiments aimed at evaluating the suitability of an easily transformable *Pichia (Ogataea) angusta* haploid strain to identify biocontrol-minus mutant clones: while the wild-type strain proved effective in reducing brown rot lesion caused by *Monilinia fructicola* on apple fruit, its derivative leucine-auxotrophic mutant L1 had no significant effect in controlling the pathogen. The addition of exogenous leucine fully restored the biocontrol capability of mutant L1, whereas a leucine stand-alone treatment showed no significant biocontrol effect (Fiori et al. 2008).

Biofilm formation may also be considered a specific and highly successful strategy to compete for space. Biofilms are microbial communities that live and grow on surfaces and can be comprised of a single species or represent multi-species consortia (Costa-Orlandi et al. 2017). Biofilms may exhibit vastly different properties as compared to free-floating cells and are considered a virulence factor for pathogenic microbes (Costa-Orlandi et al. 2017; Davey and O'Toole 2000; Desai et al. 2014). The development of a yeast biofilm starts with the adhesion of individual cells to a surface and usually involves cell wall modifications, secretion of an extracellular matrix, and often the formation of hyphae or pseudohyphae (Cavalheiro and Teixeira 2018; Costa-Orlandi et al. 2017). The process has been studied in detail and at the molecular level in medically relevant and model yeasts (Cabral et al. 2014; Cavalheiro and Teixeira 2018; d'Enfert and Janbon 2016; Lohse et al. 2018; Reynolds and Fink 2001). In biocontrol yeasts, biofilm formation, mainly in the phyllo- and carposphere (i.e., in wounds), is now considered an important mode of action and has been widely studied. However, the molecular underpinnings of the process and the composition of different biofilms (e.g., cell differentiation, multispecies biofilms) have only been studied in detail for *Pichia fermentans*. This species proved particularly intriguing in this respect, because biofilm formation in apple wounds protects against postharvest diseases, while on peaches *P. fermentans* switches from the yeast-like to the hyphal growth form and causes rapid decay of inoculated fruits in the absence of a plant pathogen (Fiori et al. 2012; Giobbe et al. 2007; Maserti et al. 2015; Sanna et al. 2012, 2013). Based on this "Jekyll & Hyde" pathogenic behaviour of *P. fermentans* on peach fruit, the capability to differentiate hyphae and pseudohyphae under particular growth conditions (e.g., depending on the nitrogen source) has been proposed as a potential biohazard factor for biocontrol yeasts (Giobbe et al. 2007). Besides *P. fermentans*, biofilm formation has also been implicated in the protective




Fig. 2 Colonisation **a** of the inner surface of an apple wound by the *Saccharomyces cerevisiae flor* strain M25. **b** *Penicillium expansum* germ tubes grow onto the yeast cells, but contact with the apple tissue

and biocontrol activities of *A. pullulans*, *Kloeckera apiculata*, *S. cerevisiae*, *Pichia kudriavzevii*, *W. anomalus*, and *M. pulcherrima* (Supplementary Table 1) (Chi et al. 2015; Klein and Kupper 2018; Ortu et al. 2005; Pu et al. 2014; Wachowska et al. 2016). In a *S. cerevisiae flor* strain, biofilm cells were far more efficient than planktonic cells in colonising the inner surface of apple wounds, thereby controlling the development of blue mould caused by *P. expansum* (Ortu et al. 2005; Scherm et al. 2003) (Fig. 2).

Secreted enzymes

The secretion of enzymes degrading cellular components is a common feature in all kinds of host–pathogen interactions and has been intensively studied. Usually, such enzymes are upregulated in nutrient poor conditions and serve the provision of nutrients (e.g., carbon sources, amino acids) that are released from “prey” cells, which may lead to the killing of these cells (i.e., mycoparasitism/fungivory; see below). Secreted enzymes such as chitinases, glucanases, or proteases are thus regularly reported and highlighted in antagonistic yeasts and implicated in their biocontrol activity.

Chitinases

The secretion of chitinolytic enzymes is considered a desirable characteristic for biocontrol agents as it allows degrading fungal cell walls (Zajc et al. 2019). Chitin degrading activity has been measured in biocontrol yeasts of the genera *Aureobasidium*, *Candida*, *Debaryomyces*, *Metschnikowia*, *Meyerozyma*, *Pichia*, *Saccharomyces*, *Tilletiopsis*, and *Wickerhamomyces* and in *Saccharomycopsis*, chitinase expression

is prevented by a thick yeast cell layer. The presence of an extracellular matrix is likely to assure an effective protection of the apple tissue (Ortu et al. unpublished)

was detected in the presence of prey cells (Supplementary Table 1) (Bar-Shimon et al. 2004; Junker et al. 2019; Lopes et al. 2015; Pretschner et al. 2018; Saravanakumar et al. 2009; Urquhart and Punja 2002; Zajc et al. 2019; Zhang et al. 2011). So far, the corresponding chitinase-encoding genes have not been cloned, deleted or overexpressed to unequivocally link these enzymes to biocontrol activity. However, chitinases from other sources than yeasts (i.e., filamentous fungi and bacteria) have demonstrated biocontrol activity against plant pathogenic fungi and chitinases are widely studied as potential biopesticides, targets for resistance breeding, or as transgenes in genetically modified plants (Dahiya et al. 2006; Herrera-Estrella and Chet 1999; Nagpure et al. 2014). Chitinases likely also affect biocontrol activity indirectly, because chito-oligosaccharides (CHOS; the results of chitin degradation) are potent inducers of plant immune responses (Kombrink et al. 2011; Langner and Gohre 2015; Liu et al. 2012, 2014).

Glucanases

Glucans are major cell wall components in fungi and exoglucanases are involved in cell wall modification, cell adhesion, and killer toxin resistance (Adams 2004; Jiang et al. 1995; Tsai et al. 2011; Xu et al. 2013). A 1,3- β -glucanase (*CoEXG1*) from *Candida oleophila* was the first gene cloned in this organism (Segal et al. 2002). Initial overexpression or deletion analyses of *CoEXG1* did not significantly affect spore germination of *Penicillium digitatum* (Yehuda et al. 2003), but later studies documented reduced inhibitory activity of the β -exoglucanase deletion mutant as compared to the wild type and overexpressing strain (*in vitro* and *in*

fructo), thereby proving the involvement of glucanases in yeast biocontrol activity (Bar-Shimon et al. 2004). In *Wickerhamomyces anomalus* (*Pichia anomala*), the deletion of the two exo- β -glucanases (*PaEXG1* and *PaEXG2*) significantly reduced biocontrol activity on fruits against *Botrytis cinerea* (Friel et al. 2007), while the single deletion of *PaEXG2* did not reduce biocontrol performance (Grevesse et al. 2003). At the transcriptional level, differential upregulation of two *W. anomalus* (*P. anomala*) exoglucanase genes was shown during the interaction with plant pathogenic fungi on infected fruits or growth with fungal cell wall preparations (Parafati et al. 2017a). Exoglucanase activity has also been detected in numerous biocontrol yeasts (see Supplementary Table 1) (Chan and Tian 2005) and was linked to antagonistic activity, but without demonstrating a causal involvement. In *Rhodotorula glutinis* and *Cryptococcus laurentii*, β -1,3-glucanase activity did not correlate with the respective inhibitory activity against *B. cinerea* (Castoria et al. 1997). Six *S. cerevisiae* isolates exhibiting antifungal activity against *Colletotrichum acutatum* secreted exoglucanases, as did a *Pichia guilliermondii* biocontrol isolate (Lopes et al. 2015; Zhang et al. 2011).

Lipases

Lipolytic activity is frequently found when screening for extracellular enzymatic activity in yeast and yeast-like strains (Arroyo-Lopez et al. 2008; Buzzini and Martini 2002; Hernandez et al. 2007). This trait has been related to the consumption of previously accumulated lipids (in the so-called 'oleaginous' yeasts), and to cold tolerance in extremophilic yeasts (Białkowska and Turkiewicz 2014; Breuer and Harms 2006; Papanikolaou and Aggelis 2011; Szczęsna-Antczak et al. 2014). Besides this, lipase activity has been detected and shown to be involved in the pathogenicity of yeasts such as *Candida*, *Cryptococcus*, or *Malassezia* species (Mayer et al. 2013; Park et al. 2013; Sommer et al. 2016). Since a number of studies have highlighted the role of lipases in the biocontrol efficacy of bacteria and fungi against plant diseases and pests (Ali et al. 2009; Berto et al. 2001; Beys da Silva et al. 2010a, b; Keyhani 2018; Manuel et al. 2012; Sánchez-Pérez et al. 2014; Vial et al. 2007; Zha et al. 2014), the lipolytic activity of antagonistic yeasts may represent a promising target for innovative studies on biological control applications.

Proteases

Although proteases are important virulence factors in entomopathogenic fungi and filamentous mycoparasites, they have been scarcely studied in biocontrol yeasts (Supplementary Table 1). Since protease activity was only detected at later growth stages (after 6–8 days of growth in

nutrient rich medium) in *C. oleophila* cultures, a minor function in biocontrol activity was hypothesised (Bar-Shimon et al. 2004). In contrast, the alkaline serine protease Alp5 from *A. pullulans* reduced spore germination and germ tube length of *Penicillium expansum*, *B. cinerea*, *M. fructicola* and *Alternaria alternata* *in vitro* and exhibited a concentration-dependent inhibitory effect on these pathogens on apple (Banani et al. 2014; Zhang et al. 2012). Protease activity has also been reported in the genera *Metschnikowia*, *Pichia*, and *Wickerhamomyces*, but not further studied or confirmed (Pretscher et al. 2018). Finally, *Saccharomycopsis* protease (and also glucanase) transcripts were significantly enriched during predation, but neither functionally investigated (Juncker et al. 2019).

Toxin production

Yeasts are not known as prolific producers of secondary metabolites, which is one of the reasons why they often raise less biosafety concerns. Consequently, relatively few toxic molecules that may contribute to biocontrol activity have been described (Supplementary Table 1). Flocculosin is a low molecular weight cellobiose lipid produced by the biocontrol yeast *Pseudozyma flocculosa* (Mimee et al. 2005, 2009; Teichmann et al. 2011). *A. pullulans* (introduced in more detail below) produces diverse polymers (e.g., pullulan, aubasidan-like exopolysaccharide, poly(β -L-malic acid)), lipids, volatiles, enzymes, and secondary metabolites. Some of these metabolites (e.g., aureobasidins, liamocins, 2-propylacrylic acid, 2-methylenesuccinic acid) confer antagonistic activity against bacteria or fungi (Prasongsuk et al. 2018; Price et al. 2013, 2017; Takesako et al. 1991; Zain et al. 2009). Toxin production provided a competitive advantage to *A. pullulans* under dry, oligotrophic conditions, whereas it had no effect (as compared to yeasts not producing toxins) on antagonistic activity in more humid environments (McCormack et al. 1995). The most prominent toxins produced by many biocontrol yeast strains are proteinaceous killer toxins (Supplementary Table 1) (Bajaj et al. 2013; Banjara et al. 2016; Belda et al. 2017; Buzdar et al. 2011; Buzzini et al. 2004; Chen et al. 2000; Coelho et al. 2009; Comitini and Ciani 2011; Comitini et al. 2004; da Silva et al. 2008; De Ingeniis et al. 2009; Golubev et al. 2006; Guo et al. 2013; Guyard et al. 2002a, b; Hua et al. 2010; Kasahara 1994a, b; Klassen et al. 2004; Marquina et al. 2001; Ramirez et al. 2015; Rodriguez-Cousino et al. 2011; Santos and Marquina 2004a, b; Santos et al. 2002, 2009; Suzuki and Nikkuni 1994; Vepstaite-Monstavice et al. 2018; Wang et al. 2007, 2012a; Weiler and Schmitt 2003). These proteins were originally identified in *S. cerevisiae* and seem to mainly kill competing yeast species (Luksa et al. 2015; Schmitt and Breinig 2006). Yeast killer toxins have thus been mainly studied with respect to the control of spoilage yeasts in

the beverage and food industry or for medical applications (Chessa et al. 2017; Chi et al. 2010; Lowes et al. 2000; Man-nazzu et al. 2019; Schmitt and Breinig 2002). However, several of these toxins also inhibit or kill plant pathogenic fungi and were thus proposed for plant protection (Corbaci and Ucar 2018; Liu et al. 2015; Marquina et al. 2002; Perez et al. 2016). Nevertheless, further investigations to evaluate the specificity of yeast toxins and assess their effects on other beneficial microorganisms (e.g., in the phyllosphere, in soil microbiota and, in the case of edible commodities, the human gut) are required, particularly in the light of a possible registration.

Volatile organic compounds

Volatile organic compounds (VOCs) are small (usually < 300 Da) molecules with low water solubility and high vapour pressure. VOCs include a panoply of molecular classes, including hydrocarbons, alcohols, thioalcohols, aldehydes, ketones, thioesters, cyclohexanes, heterocyclic compounds, phenols and benzene derivatives (Morath et al. 2012). The chemical composition of each blend of volatiles (the so-called volatilome) may change depending on the producing yeast, the antagonised pathogen and the ecological niche where the cross-talking species are growing (Parafati et al. 2017b). Recent experimental evidence has revealed the key role of the yeast volatilome in yeast-pathogen interactions, including postharvest pathogens, and mycotoxin-producing fungi (Supplementary Table 1) (Bruce et al. 2003; Lemos Jr 2016; Parafati et al. 2015). Volatiles produced by *A. pullulans* proved efficient in reducing the growth and infection by *B. cinerea*, *C. acutatum*, *P. expansum*, *P. digitatum* and *P. italicum* both *in vitro* and *in planta* (Di Francesco et al. 2014). The biocontrol activity of different food yeasts such as *W. anomalus*, *M. pulcherrima*, *S. cerevisiae* and *A. pullulans* against *B. cinerea* *in vitro* and on table grape berries was largely attributed to the production of VOCs (Parafati et al. 2015). Similarly, VOCs released by *C. sake* reduced the incidence of apple rot caused by *P. expansum* and *B. cinerea* (Arrarte et al. 2017). The inhibitory activity of *Sporidiobolus pararoseus* on spore germination and mycelial growth of *B. cinerea* was mainly attributed to 2-ethyl-1-hexanol (Huang et al. 2012), whereas *Candida intermedia* produced 1,3,5,7-cyclooctatetraene, 3-methyl-1-butanol, 2-nonanone, and phenylethyl alcohol as the major components of its volatilome during the interaction with this pathogen (Huang et al. 2011). VOCs released by *P. anomala*, *Pichia kluyveri*, and *Hanseniaspora uvarum* inhibited *Aspergillus ochraceus* growth and ochratoxin A production during processing of coffee (Masoud et al. 2005), and 2-phenylethanol was identified as the key component of the *P. anomala* volatilome preventing spore germination, mycelial growth and toxin production by *Aspergillus flavus*

(Hua et al. 2014). More than twenty different VOCs were identified in the volatilomes of selected biocontrol strains of *Cyberlindnera jadinii*, *Candida friedrichii*, *C. intermedia*, and *Lachancea thermotolerans*, but 2-phenylethanol was the most abundant and responsible for the inhibition of both mycelial growth and ochratoxin A production by *Aspergillus carbonarius* and *A. ochraceus* (Farbo et al. 2018; Fiori et al. 2014; Tilocca et al. 2019).

Mycoparasitism

Mycoparasitism (or fungivory, i.e., the consumption of one fungus by another) is rarely described and poorly studied in yeasts. *P. guilliermondii* was shown to strongly adhere to hyphae of the plant pathogen *B. cinerea* and to cause hyphal collapse, presumably due to the secretion of hydrolytic enzymes such as glucanases (see above) (Wisniewski et al. 1991). Similarly, the yeast-like Ustilaginomycete *Pseudozyma aphidis* parasitises the powdery mildew pathogen *Podosphaera xanthii* and *B. cinerea* (Calderon et al. 2019; Gafni et al. 2015). The genus *Saccharomycopsis*, comprising predacious yeasts directly feeding on their prey, was studied with respect to biocontrol of different *Penicillium* species as well as clinically relevant yeasts (Junker et al. 2017, 2018, 2019; Lachance and Pang 1997; Pimenta et al. 2008).

Induction of resistance

Plants feature an innate immune system that recognises and responds to the presence of microorganisms (Chisholm et al. 2006; Jones and Dangl 2006). This plant immune response can induce resistance systemically and is the basis for the application of microorganisms as plant fertilisers and fortifiers (Gozzo and Faoro 2013; Pieterse et al. 2014). Biocontrol yeasts can elicit systemic resistance of plants against a broad range of pathogens (Supplementary Table 1) (Barda et al. 2015; Buxdorf et al. 2013a, b; Lee et al. 2017; Liu et al. 2016) and this activity is suggested to contribute to their biocontrol activity. For example, *S. cerevisiae*, *Rhodospodium paludigenum*, *Candida saitoana*, *C. oleophila* and *Metschnikowia* species induce an innate immune response and eventually cause resistance against phyllosphere pathogens in fruits (De Miccolis Angelini et al. 2019; Droby et al. 2002; El Ghaouth et al. 2003; Hadwiger et al. 2015; Herzhkowitz et al. 2012; Lu et al. 2013, 2014; Sun et al. 2018). In the case of *C. oleophila*, this induction has been attributed to the overproduction of reactive oxygen species in the plant (Macarasin et al. 2010), but yeast cell components (from dead cells) can also trigger systemic resistance (De Miccolis Angelini et al. 2019). Living cells are consequently not always required for such induction. In some cases, biocontrol yeasts such as *C. laurentii*, *Cryptococcus flavescens*, and *R. glutinis* have been used in combination with resistance

inducers such as salicylic acid or rhamnolipids (Yan et al. 2014; Yu and Zheng 2006; Zhang et al. 2007c).

Registered biocontrol yeast species

There is a huge discrepancy between the plethora of “biocontrol yeasts” described in scientific publications and the few yeast-based plant protection products that are registered and marketed as plant protection products. A range of factors (e.g., lack of mechanistic understanding, hurdles/costs of registration, lack of partners/consortia with required expertise, little commercial potential) are likely responsible for this apparent difficulty to develop yeast-based plant protection products. Here, we briefly highlight five yeast species (*C. oleophila*, *A. pullulans*, *M. fructicola*, *C. albidus*, and *S. cerevisiae*) that are currently or have been registered as plant protection agents.

Candida oleophila

Species of the genus *Candida* are often isolated from environmental samples and many isolates strongly inhibit plant pathogens. Representatives are, for example, *C. diversa* (Li et al. 2016; Liu et al. 2017), *C. ernobii* (Liu et al. 2010), *C. guillemontii* (McLaughlin et al. 1992; Papon et al. 2013; Saligkari et al. 2002), *C. oleophila* (Droby et al. 2002; Gamagae et al. 2003; Lahlali et al. 2004; Molinu et al. 2011; Wang et al. 2012b), *C. saitoana* (Arras et al. 2006, 2010; El-Ghaouth et al. 1998, 2000a, b, c), *C. sake* (Arrarte et al. 2017; Calvo-Garrido et al. 2013; Canamas et al. 2008; Carbo et al. 2018; McLaughlin et al. 1992; Morales et al. 2008; Nunes et al. 2002a, b; Torres et al. 2006; Usall et al. 2000; Yehuda et al. 2003), or *C. subhashii* (Hilber-Bodmer et al. 2017) that have all been envisioned as biocontrol agents against mold and postharvest diseases of pome, stone and citrus fruit. In particular for *C. sake*, a wealth of studies on production and formulation have been performed in order to render postharvest biocontrol more reliable and efficacious (Abadias et al. 2000, 2001a, b, c, 2003; Canamas et al. 2008; Carbo et al. 2018; Nunes et al. 2002a, b; Torres et al. 2003, 2006; Usall et al. 2000).

C. oleophila was the first yeast to be developed into a commercial plant protection agent and the fundamental research accompanying this initiative has established, for the first time, different mechanisms underlying the antifungal activity of yeasts in general. Although yeasts are generally believed to antagonise plant pathogenic fungi due to their competition for nutrients and space, the work on *C. oleophila* and other *Candida* species identified hydrolytic enzymes such as proteases, chitinases and glucanases, as well as volatile compounds, that have been implicated in antifungal activity (Bar-Shimon et al. 2004; Huang et al. 2011; Segal

et al. 2002) (also see above and Supplementary Table 1). Furthermore, biofilm formation, high osmotolerance, induction of resistance in the plant/fruit, and direct parasitism of hyphae were shown to contribute to the biocontrol activity of *Candida* species (Droby and Chalutz 1994; Droby et al. 2002; El-Ghaouth et al. 2003; Wisniewski et al. 1995, 2007). To overcome the inconsistent performance of the initial *Candida*-based biocontrol products (and of early biological plant protection products in general), combinations with fungicides, different buffers (e.g., calcium chloride, bicarbonate), chitosan, or lysozyme were studied (Droby et al. 1998, 2003a, b; El-Ghaouth and Wilson 2002; Scherm et al. 2003; Wilson and El-Ghaouth 2002). *C. oleophila* was also transformed, by electroporation and with the hygromycin B gene as a marker, to study its mode of antagonism at the molecular level (Yehuda et al. 2001).

The *C. oleophila* strains I-182 and O have been developed into the biocontrol products Aspire[®] and Nexy[®], respectively. The latter was the first biocontrol yeast to be registered against a postharvest disease (Wisniewski et al. 2007) and *C. oleophila* strain O has been approved as a plant protection agent in Europe in 2013 (European Commission Health & Consumers Directorate-General 2013; European Food Safety Authority (EFSA) 2015a).

Aureobasidium pullulans

The saprophytic ascomycete *A. pullulans* is frequently isolated from leaf, flower or soil samples, occurs worldwide, and exhibits a polymorphic appearance. Biocontrol activity has been documented for several *A. pullulans* strains, but only DSM 14940 (CF 10) and DSM 14941 (CF 40) are registered, in mixture, as active ingredients of plant protection products against the fireblight disease caused by the bacterium *Erwinia amylovora* and postharvest diseases (European Food Safety Authority (EFSA) 2013). These two *A. pullulans* strains were selected based on their strong inhibition towards *E. amylovora* in co-culture experiments at high synthetic nectar concentration (25%) (Seibold et al. 2004). The two isolates DSM 14940 (CF 10) and DSM 14941 (CF 40) also exhibited stronger inhibitory activity, in detached flower assays, than other bacterial and yeast antagonists (Kunz 2004). The two strains were formulated as a wettable powder under the product name Blossom-Protect[®] and tested under field conditions at different sites and over several years (Kunz 2004; Kunz and Haug 2006; Kunz et al. 2011; Seibold et al. 2004). The same two *A. pullulans* strains were also developed and registered to control postharvest diseases of apple as the product Boni-Protect[®] (Weiss and Mögel 2006). Similar applications against storage and rot diseases of strawberries, plum and sour cherries are being studied (Holb and Kunz 2013; Weiss et al. 2014).

The two registered products containing *A. pullulans* as an active ingredient are interesting from different points of view. Contrary to the large majority of biocontrol products, Blossom- and Boni-Protect[®] contain two different strains, albeit belonging to the same species. As for many other biocontrol yeasts, the *A. pullulans* mode of action involves competition for space and nutrients, but enzymes such as proteases, chitinases or secreted molecules (see above) may also be involved. Specific metabolites or enzymes and their contribution to the biocontrol activity of DSM 14940 (CF 10) and DSM 14941 (CF 40) have not been identified and the strains do not seem to have been characterised genetically. In contrast to most registered plant protection products, including biological products, the original Blossom-Protect[®] has a rather limited range of application. The expansion to novel indications, beyond fireblight of pome fruit trees, is thus certainly also motivated by economic needs.

Metschnikowia fructicola

The genus *Metschnikowia* comprises species of mainly phyllosphere and nectar yeasts that are globally distributed (Chappell and Fukami 2018; Lachance et al. 2001; Pozo et al. 2012; Slavikova et al. 2007; Vadkertiova et al. 2012). Among those, *M. fructicola* and *M. pulcherrima* are the most studied with respect to biocontrol, are able to inhibit a range of postharvest and plant rot diseases, and include the most potent antagonistic yeasts that have ever been identified (Akgun Karabulut et al. 2003; Hilber-Bodmer et al. 2017; Parafati et al. 2015; Piano et al. 1997; Saravanakumar et al. 2008; Spadaro et al. 2010a, b; Turkel et al. 2014). Complete genomes are available for several *Metschnikowia* species, including *M. fructicola* and *M. pulcherrima* (Gore-Lloyd et al. 2019; Piombo et al. 2018), and transformation protocols have been established and used to express green fluorescent protein and complement a naturally occurring mutant (Gore-Lloyd et al. 2019; Nigro et al. 1999).

The strong antifungal activity of *Metschnikowia* species is mediated by a range of mechanisms that involve competition for nutrients (e.g.; amino acids, iron), secretion of glucanases and chitinases, and the production of volatile organic compounds (Banani et al. 2015; Dhami et al. 2016; Gore-Lloyd et al. 2019; Hershkovitz et al. 2013; Saravanakumar et al. 2008; Sipiczki 2006; Zajc et al. 2019). The application of *Metschnikowia* cells to fruits (e.g., grapefruit) also induces an oxidative burst in the plant tissue that eventually results in the activation of plant defense responses (Hershkovitz et al. 2012; Macarasin et al. 2010).

Originally, *M. fructicola*, isolate NRRL Y-30752, was isolated and discovered in Israel and developed and registered as a biocontrol product for preventing postharvest diseases, particularly in sweet potato and carrot (Eshel et al. 2009; Kurtzman and Droby 2001; Wisniewski and Droby

2012). *M. fructicola* has also been patented as an antagonist of plant pathogenic microorganisms (Droby and El-Gerberia 2006). It seems that over time different companies showed interest in developing a biological fungicide based on *M. fructicola* NRRL Y-30752, but the isolate is now pursued by Koppert Biological Systems and has recently been approved as a plant protection agent against fungal diseases in stone fruits, strawberries and grapes by the European Food Safety Authority (EFSA) (European Food Safety Authority (EFSA) 2015c, 2017).

Cryptococcus albidus

Basidiomycetes of the genus *Cryptococcus* are widespread in nature and frequently isolated from water sources, soil and decaying plant material. Studied for their potential to produce high lipid yields for biodiesel production, strains of *C. albidus*, *C. laurentii* and *C. flavus* have also been shown to protect peach, cherry, strawberry, tomato, citrus and pome fruits against postharvest decay (Elad et al. 1994; Tian et al. 2004; Zhang et al. 2007a, b). *C. albidus* was used as a biocontrol agent in the product Yieldplus[®], which was registered in 1997 and marketed by Anchor Bio-Technologies in South Africa. This product was sold for over 15 years, but it has now been withdrawn from the market (Mbili 2012). Yieldplus[®] was formulated for pome and citrus fruits against *B. cinerea* and *P. expansum* and later shown to be effective in the control of *Botrytis* during post-harvest cold storage of strawberries (Kowalska et al. 2012).

Regarding the mode of antagonism, most of the evidence points to competition for nutrients and space. Culture filtrates do not show any inhibitory activity against *B. cinerea* or *P. expansum*. However, both pathogens show reduced conidial germination and germ tube growth in liquid co-cultures (Fan and Tian 2001; Helbig 2002). The addition of glucose or NH₄NO₃ to the medium reduces biocontrol ability against *P. expansum*, but not against *B. cinerea* (Lutz et al. 2013). Beside nutrient competition, little conclusive evidence is available to determine a mode of antagonism. *C. albidus* exhibits glucanase, chitinase and protease activity in the corresponding substrate media. It also produces unidentified volatile compounds that inhibit fungal growth and can display killer activity against *C. glabrata* (Lutz et al. 2013). However, none of these mechanisms have been directly linked to the inhibitory activity of the target plant pathogens.

Saccharomyces cerevisiae

S. cerevisiae is mainly known as a model organism for cell biology, its biotechnological usage, and most importantly the application in food and beverage production. Envisioning *S. cerevisiae* for biocontrol may be motivated by its perception as a safe organism that can be more easily registered,

but also by its model organism status and the feasibility of molecular analyses. Overall, the model *S. cerevisiae* isolate BY4741 exhibited an intermediate antifungal activity against filamentous fungi and in comparison to a broad collection of wild yeast isolates (Hilber-Bodmer et al. 2017). This laboratory strain is thus ideally suited as a model host to express genes potentially involved in biocontrol activity and thereby improving or weakening its antifungal action.

A number of *S. cerevisiae* strains (e.g., DISAABA1182, RC008, RC009, RC012, and RC016) reduced the growth of plant pathogens such as *A. carbonarius*, *A. ochraceus*, *A. parasiticus* or *Fusarium graminearum* and also inhibited mycotoxin (e.g., aflatoxin, ochratoxin A, zearalenone, deoxynivalenol) production by these species (Armando et al. 2012a, 2013; Cubaiu et al. 2012). The mycotoxin-removing activity is due to adsorption to *S. cerevisiae* cell walls, stress responses to the toxin (e.g., changes in plasma membrane composition following patulin exposure), as well as direct transcriptional downregulation of polyketide synthesis (Armando et al. 2012b; Cubaiu et al. 2012; Oporto et al. 2019). Other biocontrol mechanisms employed by *S. cerevisiae* include the secretion of killer activity and hydrolytic enzymes as well as organic volatile compounds in yeasts that have been described and studied with respect to their antifungal activity against *C. acutatum* on citrus (Lopes et al. 2015) (also see Supplementary Table 1). Volatiles in general and specific compounds (e.g., alcohols and esters), were also identified in *S. cerevisiae* and implicated in the biocontrol activity against the citrus black spot disease caused by *Guignardia citricarpa* or postharvest decay in strawberries (Fialho et al. 2010; Oro et al. 2017). Seed, soil or foliar applications of a dried, active *S. cerevisiae* preparation also had a plant growth promoting effect and showed biocontrol activity against soilborne fungal pathogens such as *Fusarium*, *Sclerotium* or *Rhizoctonia* (Shalaby and El-Nady 2008). Finally, comprehensive transcriptome analyses have confirmed that the application of a cell wall preparation of the *S. cerevisiae* strain LAS117 (i.e., cerevisane[®]) induces the expression of genes involved in the plant response to fungal attack (De Miccolis Angelini et al. 2019). *S. cerevisiae* is therefore considered a promising biocontrol and probiotic organism for reducing growth of fungal pathogens and mycotoxins in fruit, vegetable and feedstuff (Cubaiu et al. 2012; Dogi et al. 2011; Pizzolitto et al. 2012; Prado et al. 2011). However, the only registered, active compound and commercial biocontrol application of Brewer's yeast (*S. cerevisiae*) is the product Romeo[®] with cerevisane[®] as the active ingredient (European Food Safety Authority (EFSA) 2015b). This preparation is used as a preventive inducer of systemic resistance against powdery and downy mildew in grapes, fruits and vegetables and thus represents an application and plant protection product that differs from other such solutions insofar it is not based on active, living cells.

Conclusions and outlook

The disparity between the number of yeast species exerting biocontrol activity against specific plant pathogens in laboratory assays and the number of yeast that are actually registered and successfully employed as plant protection products is likely caused by the lack of mechanistic understanding, the costs of registration, the lack of partners/consortia with required expertise, or a limited commercial potential. However, the general trend towards reduced pesticide use will certainly favour and create more incentives for the development of alternative plant protection solutions such as biocontrol yeasts. In the environment, these organisms interact intraspecifically, as well as with other microbes (including, but not exclusively, with plant pathogens) and host plants. These complex interactions and interdependencies eventually determine whether a disease sets in or an antagonistic yeast suppresses a pathogen and supports plant health. It is impossible to manage—for plant protection applications—such complex interactions without detailed knowledge of the interacting bionts. By studying and identifying modes of action in the laboratory, a reductionist approach is thus an important first step in the development and successful application of biocontrol in general.

Nowadays, the breakthrough achievements in the field of system biology, molecular biology and the related computational tools enable revealing the structural and functional peculiarities of any potential biocontrol agent. Exploiting these tools for the investigation and prediction of functional dynamics occurring between antagonists opens new avenues for the design of consortia of microbial antagonists that synergistically cooperate for the biocontrol of plant pathogens. Although strain mixtures for the biocontrol of plant pathogens are already available commercially (e.g., Blossom- and Boni-Protect[®], see above) and described in the literature (Heydari and Pessaraki 2010; Lopes et al. 2012; Spadaro and Gullino 2005), employment of taxonomically divergent, but functionally complementary strains might represent a promising approach to follow in the near future in an attempt to design a standardised, multi-targeted, efficacious biocontrol strategy.

A deep mechanistic insight does not guarantee a successful product development and registration, but we argue that fundamental research on biocontrol mechanisms is a key aspect for successful biocontrol applications and at the same time still a frontier in biocontrol research. Hardly any antagonistic mechanism employed by biocontrol yeasts is understood and unequivocally proven by gene deletions and overexpression. This lack of fundamental understanding is also one of the reasons why, so far, little efforts were undertaken to improve, either by selection or molecular tools, biocontrol yeasts for plant protection. In general, molecular tools (e.g.,

gene deletion or overexpression, introduction of trans-genes, synthetic biology techniques) were rarely used in biocontrol yeasts, even though these technologies have the potential to empower studying and understanding these organisms at a whole new level (Marchand et al. 2007). Probably the first “biocontrol engineering example” among yeasts is a *S. cerevisiae* strain expressing the antifungal peptide cecropin A, which resulted in complete inhibition of *Colletotrichum coccoodes* growth on tomatoes (Jones and Prusky 2002). Cecropin A was also expressed in *Pichia pastoris*, generating a strain controlling apple blue mold caused by *P. expansum* (Ren et al. 2012). A *P. pastoris* strain was also engineered for improved control of *P. expansum* by expressing the recombinant peach and pea defensins rDFN1 or rPsd1, respectively (Janisiewicz et al. 2008; Wisniewski et al. 2003). Besides this, only killer toxin activity has been extensively studied and transferred to new strains; mainly for biotechnological applications (Bajaj and Sharma 2010; Bussey et al. 1988; Schmitt and Schernikau 1997). Although the current legislation forbids the deliberate release of genetically manipulated microbials in the environment, model yeasts could also be engineered to shed light on the molecular mechanisms governing their antagonistic capability, their persistence on the host plant, or to better understand how to limit their capability to spread and interbreed (Callaway 2018; Goold et al. 2018; Klemsdal and Tronsmo 1999; Maselko et al. 2017).

Another frontier for biocontrol yeast research is the application in soil, for the protection against soilborne plant pathogens. Phyllosphere applications, particularly the use of yeasts against postharvest diseases, have been identified as the most promising target (Wisniewski et al. 2007). However, a broad range of yeast species occur predominantly in soil, exhibit strong antifungal activity, and could be envisioned as plant protection agents in this environment (Botha 2011; Hilber-Bodmer et al. 2017; Yurkov 2018). Developing yeasts as biocontrol agents against soilborne plant diseases is also attractive, because only few control options are available for the many severe soilborne plant diseases.

The development of a (yeast) biocontrol product depends on a chain of activities and disciplines, from science to industry and legislation, which have to come together, interact and build upon one another. Strengthening the exchange and interaction among these disciplines is thus essential to foster the commercialisation of biocontrol products (Usall et al. 2016). However, establishing such a virtuous cycle is difficult because of the different interests and qualities required and can be further hampered by economic constraints, commercial interests (and thereby a hesitation to share know-how or even material), or a lack of actors with complementary expertise. Considering the fact that many biocontrol solutions are local endeavours, have a limited potential to incur financial gains (in particular in comparison to medical applications), and are somehow idealistic in

nature, commercial interests may actually rather harm than benefit the development of commercial biocontrol solutions. In particular, governmental research institutions, engaging in fundamental and applied research and being less driven by commercialisation, may take up a crucial spot for developing biocontrol solutions in the future.

Acknowledgements FMF acknowledges funding from the Swiss National Science Foundation (SNSF, Grant 31003A_175665/1). This paper is dedicated to the memory of Enrico Berardi, sailor, dreamer, and yeast geneticist.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abadias M, Teixido N, Usall J, Viñas I, Magan N (2000) Solute stresses affect growth patterns, endogenous water potentials and accumulation of sugars and sugar alcohols in cells of the biocontrol yeast *Candida sake*. *J Appl Microbiol* 89:1009–1017. <https://doi.org/10.1046/j.1365-2672.2000.01207.x>
- Abadias M, Benabarre A, Teixido N, Usall J, Viñas I (2001a) Effect of freeze drying and protectants on viability of the biocontrol yeast *Candida sake*. *Int J Food Microbiol* 65:173–182. [https://doi.org/10.1016/S0168-1605\(00\)00513-4](https://doi.org/10.1016/S0168-1605(00)00513-4)
- Abadias M, Teixido N, Usall J, Benabarre A, Viñas I (2001b) Viability, efficacy, and storage stability of freeze-dried biocontrol agent *Candida sake* using different protective and rehydration media. *J Food Prot* 64:856–861. <https://doi.org/10.4315/0362-028X-64.6.856>
- Abadias M, Teixido N, Usall J, Viñas I, Magan N (2001c) Improving water stress tolerance of the biocontrol yeast *Candida sake* grown in molasses-based media by physiological manipulation. *Can J Microbiol* 47:123–129
- Abadias M, Teixido N, Usall J, Viñas I (2003) Optimization of growth conditions of the postharvest biocontrol agent *Candida sake* CPA-1 in a lab-scale fermenter. *J Appl Microbiol* 95:301–309
- Adams DJ (2004) Fungal cell wall chitinases and glucanases. *Microbiology* 150:2029–2035. <https://doi.org/10.1099/mic.0.26980-0>
- Akgun Karabulut O, Smilanick J, Mlikota Gabler F, Mansour M, Droby S (2003) Near-harvest applications of *Metschnikowia fructicola*, ethanol, and sodium bicarbonate to control postharvest diseases of grape in central California. *Plant Dis* 87:1384–1389. <https://doi.org/10.1094/PDIS.2003.87.11.1384>
- Ali S, Huang Z, Ren SX (2009) Production and extraction of extracellular lipase from the entomopathogenic fungus *Isaria fumosoroseus* (Cordycipitaceae; Hypocreales). *Biocontrol Sci Technol* 19:81–89. <https://doi.org/10.1080/09583150802588524>
- Armando MR, Dogi CA, Rosa CA, Dalcerro AM, Cavaglieri LR (2012a) *Saccharomyces cerevisiae* strains and the reduction of *Aspergillus parasiticus* growth and aflatoxin B1 production at different interacting environmental conditions, *in vitro*. *Food Addit Contam Part A* 29:1443–1449. <https://doi.org/10.1080/19440049.2012.698655>
- Armando MR et al (2012b) Adsorption of ochratoxin A and zearalenone by potential probiotic *Saccharomyces cerevisiae* strains and

- its relation with cell wall thickness. *J Appl Microbiol* 113:256–264. <https://doi.org/10.1111/j.1365-2672.2012.05331.x>
- Armando MR, Dogi CA, Poloni V, Rosa CA, Dalcerio AM, Cavaglieri LR (2013) *In vitro* study on the effect of *Saccharomyces cerevisiae* strains on growth and mycotoxin production by *Aspergillus carbonarius* and *Fusarium graminearum*. *Int J Food Microbiol* 161:182–188. <https://doi.org/10.1016/j.ijfoodmicro.2012.11.016>
- Arrarte E, Garmendia G, Rossini C, Wisniewski M, Vero S (2017) Volatile organic compounds produced by Antarctic strains of *Candida sake* play a role in the control of postharvest pathogens of apples. *Biol Control* 109:14–20. <https://doi.org/10.1016/j.biocontrol.2017.03.002>
- Arras G, Molinu MG, Dore A, Venditti T, Fois M, Petretto A, D'Hallewin G (2006) Inhibitory activity of 2-deoxy-D-glucose and *Candida saitoana* against *Penicillium digitatum*. *Commun Agric Appl Biol Sci* 71:929–936
- Arras G et al (2010) Synergic interactions between 2-deoxy-D-glucose and *Candida saitoana* enhances citrus green mould control. *Commun Agric Appl Biol Sci* 75:555–562
- Arroyo-Lopez FN, Querol A, Bautista-Gallego J, Garrido-Fernandez A (2008) Role of yeasts in table olive production. *Int J Food Microbiol* 128:189–196. <https://doi.org/10.1016/j.ijfoodmicro.2008.08.018>
- Bajaj BK, Sharma S (2010) Construction of killer industrial yeast *Saccharomyces cerevisiae* hau-1 and its fermentation performance. *Braz J Microbiol* 41:477–485. <https://doi.org/10.1590/S1517-838220100002000030>
- Bajaj BK, Raina S, Singh S (2013) Killer toxin from a novel killer yeast *Pichia kudriavzevii* RY55 with idiosyncratic antibacterial activity. *J Basic Microbiol* 53:645–656. <https://doi.org/10.1002/jobm.201200187>
- Banani H, Spadaro D, Zhang D, Matic S, Garibaldi A, Gullino ML (2014) Biocontrol activity of an alkaline serine protease from *Aureobasidium pullulans* expressed in *Pichia pastoris* against four postharvest pathogens on apple. *Int J Food Microbiol* 182–183:1–8. <https://doi.org/10.1016/j.ijfoodmicro.2014.05.001>
- Banani H, Spadaro D, Zhang D, Matic S, Garibaldi A, Gullino ML (2015) Postharvest application of a novel chitinase cloned from *Meischnikowia fructicola* and overexpressed in *Pichia pastoris* to control brown rot of peaches. *Int J Food Microbiol* 199:54–61. <https://doi.org/10.1016/j.ijfoodmicro.2015.01.002>
- Banjara N, Nickerson KW, Suhr MJ, Hallen-Adams HE (2016) Killer toxin from several food-derived *Debaryomyces hansenii* strains effective against pathogenic *Candida* yeasts. *Int J Food Microbiol* 222:23–29. <https://doi.org/10.1016/j.ijfoodmicro.2016.01.016>
- Barber MF, Elde NC (2015) Buried treasure: evolutionary perspectives on microbial iron piracy. *Trends Genet* 31:627–636. <https://doi.org/10.1016/j.tig.2015.09.001>
- Barda O, Shalev O, Alster S, Buxdorf K, Gafni A, Levy M (2015) *Pseudozyma aphidis* induces salicylic-acid-independent resistance to *Clavibacter michiganensis* in tomato plants. *Plant Dis* 99:621–626. <https://doi.org/10.1094/PDIS-04-14-0377-RE>
- Bar-Shimon M et al (2004) Characterization of extracellular lytic enzymes produced by the yeast biocontrol agent *Candida oleophila*. *Curr Genet* 45:140–148. <https://doi.org/10.1007/s00294-003-0471-7>
- Bekatorou A, Psarianos C, Koutinas AA (2006) Production of food grade yeasts. *Food Technol Biotechnol* 44:407–415
- Belda I, Ruiz J, Alonso A, Marquina D, Santos A (2017) The biology of *Pichia membranifaciens* killer toxins. *Toxins* (Basel). <https://doi.org/10.3390/toxins9040112>
- Berto P, Jijakli MH, Lepoivre P (2001) Possible role of colonization and cell wall-degrading enzymes in the differential ability of three *Ulocladium atrum* strains to control *Botrytis cinerea* on necrotic strawberry leaves. *Phytopathology* 91:1030–1036. <https://doi.org/10.1094/PHTO.2001.91.11.1030>
- Beys da Silva WO, Santi L, Correa AP, Silva LA, Bresciani FR, Schrank A, Vainstein MH (2010a) The entomopathogen *Metarhizium anisopliae* can modulate the secretion of lipolytic enzymes in response to different substrates including components of arthropod cuticle. *Fungal Biol* 114:911–916. <https://doi.org/10.1016/j.funbio.2010.08.007>
- Beys da Silva WO, Santi L, Schrank A, Vainstein MH (2010b) *Metarhizium anisopliae* lipolytic activity plays a pivotal role in *Rhipicephalus* (Boophilus) microplus infection. *Fungal Biol* 114:10–15. <https://doi.org/10.1016/j.mycres.2009.08.003>
- Białkowska A, Turkiewicz M (2014) Miscellaneous cold-active yeast enzymes of industrial importance. In: Buzzini P, Margesin R (eds) Cold-adapted yeasts: biodiversity, adaptation strategies and biotechnological significance. Springer, Berlin, pp 377–395. https://doi.org/10.1007/978-3-642-39681-6_17
- Botha A (2011) The importance and ecology of yeasts in soil. *Soil Biol Biochem* 43:1–8. <https://doi.org/10.1016/j.soilbio.2010.10.001>
- Breuer U, Harms H (2006) *Debaryomyces hansenii*—an extremophilic yeast with biotechnological potential. *Yeast* 23:415–437. <https://doi.org/10.1002/yea.1374>
- Bruce A, Stewart D, Verrall S, Wheatley RE (2003) Effect of volatiles from bacteria and yeast on the growth and pigmentation of sapstain fungi. *Int Biodeterior Biodegrad* 51:101–108. [https://doi.org/10.1016/S0964-8305\(02\)00088-4](https://doi.org/10.1016/S0964-8305(02)00088-4)
- Bussey H, Vernet T, Sdicu AM (1988) Mutual antagonism among killer yeasts: competition between K1 and K2 killers and a novel cDNA-based K1-K2 killer strain of *Saccharomyces cerevisiae*. *Can J Microbiol* 34:38–44
- Butler G et al (2009) Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* 459:657–662. <https://doi.org/10.1038/nature08064>
- Buxdorf K, Rahat I, Gafni A, Levy M (2013a) The epiphytic fungus *Pseudozyma aphidis* induces jasmonic acid- and salicylic acid/nonexpressor of PR1-independent local and systemic resistance. *Plant Physiol* 161:2014–2022. <https://doi.org/10.1104/pp.112.212969>
- Buxdorf K, Rahat I, Levy M (2013b) *Pseudozyma aphidis* induces ethylene-independent resistance in plants. *Plant Signal Behav* 8:e26273. <https://doi.org/10.4161/psb.26273>
- Buzdar MA, Chi Z, Wang Q, Hua MX, Chi ZM (2011) Production, purification, and characterization of a novel killer toxin from *Kluyveromyces siamensis* against a pathogenic yeast in crab. *Appl Microbiol Biotechnol* 91:1571–1579. <https://doi.org/10.1007/s00253-011-3220-8>
- Buzzini P, Martini A (2002) Extracellular enzymatic activity profiles in yeast and yeast-like strains isolated from tropical environments. *J Appl Microbiol* 93:1020–1025. <https://doi.org/10.1046/j.1365-2672.2002.01783.x>
- Buzzini P, Corazzi L, Turchetti B, Buratta M, Martini A (2004) Characterization of the *in vitro* antimycotic activity of a novel killer protein from *Williopsis saturnus* DBVPG 4561 against emerging pathogenic yeasts. *FEMS Microbiol Lett* 238:359–365. <https://doi.org/10.1016/j.femsle.2004.07.060>
- Cabral V et al (2014) Targeted changes of the cell wall proteome influence *Candida albicans* ability to form single- and multi-strain biofilms. *PLoS Pathog* 10:e1004542. <https://doi.org/10.1371/journal.ppat.1004542>
- Calderon CE, Rotem N, Harris R, Vela-Corcia D, Levy M (2019) *Pseudozyma aphidis* activates reactive oxygen species production, programmed cell death and morphological alterations in the necrotrophic fungus *Botrytis cinerea*. *Mol Plant Pathol* 20:562–574. <https://doi.org/10.1111/mp.12775>
- Callaway E (2018) Synthetic species made to shun sex with wild organisms. *Nature* 553:259–260. <https://doi.org/10.1038/d41586-018-00625-1>

- Calvo-Garrido C, Viñas I, Elmer P, Usall J, Teixido N (2013) *Candida sake* CPA-1 and other biologically based products as potential control strategies to reduce sour rot of grapes. *Lett Appl Microbiol* 57:356–361. <https://doi.org/10.1111/lam.12121>
- Canamas TP, Viñas I, Usall J, Magan N, Solsona C, Teixido N (2008) Impact of mild heat treatments on induction of thermotolerance in the biocontrol yeast *Candida sake* CPA-1 and viability after spray-drying. *J Appl Microbiol* 104:767–775. <https://doi.org/10.1111/j.1365-2672.2007.03590.x>
- Cao M, Gao M, Lopez-Garcia CL, Wu Y, Seetharam AS, Severin AJ, Shao Z (2017) Centromeric DNA facilitates nonconventional yeast genetic engineering. *ACS Synth Biol* 6:1545–1553. <https://doi.org/10.1021/acssynbio.7b00046>
- Carbo A, Torres R, Teixido N, Usall J, Medina A, Magan N (2018) Impact of climate change environmental conditions on the resilience of different formulations of the biocontrol agent *Candida sake* CPA-1 on grapes. *Lett Appl Microbiol*. <https://doi.org/10.1111/lam.12889>
- Castoria R, De Curtis F, Lima G, De Cicco V (1997) β -1,3-glucanase activity of two saprophytic yeasts and possible mode of action as biocontrol agents against postharvest diseases. *Postharvest Biol Technol* 12:293–300. [https://doi.org/10.1016/S0925-5214\(97\)00061-6](https://doi.org/10.1016/S0925-5214(97)00061-6)
- Cavalheiro M, Teixeira MC (2018) *Candida* biofilms: threats, challenges, and promising strategies. *Front Med (Lausanne)* 5:28. <https://doi.org/10.3389/fmed.2018.00028>
- Chan Z, Tian S (2005) Interaction of antagonistic yeasts against postharvest pathogens of apple fruit and possible mode of action. *Postharvest Biol Technol* 36:215–223. <https://doi.org/10.1016/j.postharvbio.2005.01.001>
- Chappell CR, Fukami T (2018) Nectar yeasts: a natural microcosm for ecology. *Yeast* 35:417–423. <https://doi.org/10.1002/yea.3311>
- Chen WB, Han YF, Jong SC, Chang SC (2000) Isolation, purification, and characterization of a killer protein from *Schwanniomyces occidentalis*. *Appl Environ Microbiol* 66:5348–5352. <https://doi.org/10.1128/aem.66.12.5348-5352.2000>
- Chessa R et al (2017) Biotechnological exploitation of *Tetrapisispora phaffii* killer toxin: heterologous production in *Komagataella phaffii* (*Pichia pastoris*). *Appl Microbiol Biotechnol* 101:2931–2942. <https://doi.org/10.1007/s00253-016-8050-2>
- Chi ZM, Liu G, Zhao S, Li J, Peng Y (2010) Marine yeasts as biocontrol agents and producers of bio-products. *Appl Microbiol Biotechnol* 86:1227–1241. <https://doi.org/10.1007/s00253-010-2483-9>
- Chi M et al (2015) Increase in antioxidant enzyme activity, stress tolerance and biocontrol efficacy of *Pichia kudriavzevii* with the transition from a yeast-like to biofilm morphology. *Biol Cont* 90:113–119. <https://doi.org/10.1016/j.biocontrol.2015.06.006>
- Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006) Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124:803–814. <https://doi.org/10.1016/j.cell.2006.02.008>
- Coelho AR, Tachi M, Pagnocca FC, Nobrega GM, Hoffmann FL, Harada K, Hirooka EY (2009) Purification of *Candida guilliermondii* and *Pichia ohmeri* killer toxin as an active agent against *Penicillium expansum*. *Food Addit Contam Part A* 26:73–81. <https://doi.org/10.1080/02652030802227227>
- Comitini F, Ciani M (2011) *Kluyveromyces wickerhamii* killer toxin: purification and activity towards *Brettanomyces/Dekkera* yeasts in grape must. *FEMS Microbiol Lett* 316:77–82. <https://doi.org/10.1111/j.1574-6968.2010.02194.x>
- Comitini F, Di Pietro N, Zacchi L, Mannazzu I, Ciani M (2004) *Kluyveromyces phaffii* killer toxin active against wine spoilage yeasts: purification and characterization. *Microbiology* 150:2535–2541. <https://doi.org/10.1099/mic.0.27145-0>
- Corbaci C, Ucar FB (2018) Purification, characterization and *in vivo* biocontrol efficiency of killer toxins from *Debaryomyces hansenii* strains. *Int J Biol Macromol* 119:1077–1082. <https://doi.org/10.1016/j.ijbiomac.2018.07.121>
- Costa-Orlandi CB et al (2017) Fungal biofilms and polymicrobial diseases. *J Fungi (Basel)* 3:22. <https://doi.org/10.3390/jof3020022>
- Cubaiu L, Abbas H, Dobson AD, Budroni M, Migheli Q (2012) A *Saccharomyces cerevisiae* wine strain inhibits growth and decreases ochratoxin A biosynthesis by *Aspergillus carbonarius* and *Aspergillus ochraceus*. *Toxins (Basel)* 4:1468–1481. <https://doi.org/10.3390/toxins4121468>
- da Silva S, Calado S, Lucas C, Aguiar C (2008) Unusual properties of the halotolerant yeast *Candida nodaensis* killer toxin. *CnKT. Microbiol Res* 163:243–251. <https://doi.org/10.1016/j.micres.2007.04.002>
- Dahiya N, Tewari R, Hoondal GS (2006) Biotechnological aspects of chitinolytic enzymes: a review. *Appl Microbiol Biotechnol* 71:773–782. <https://doi.org/10.1007/s00253-005-0183-7>
- Davey ME, O'Toole GA (2000) Microbial biofilms: from ecology to molecular genetics. *Microbiol Mol Biol Rev* 64:847–867. <https://doi.org/10.1128/mmr.64.4.847-867.2000>
- De Ingeniis J, Raffaelli N, Ciani M, Mannazzu I (2009) *Pichia anomala* DBVPG 3003 secretes a ubiquitin-like protein that has antimicrobial activity. *Appl Environ Microbiol* 75:1129–1134. <https://doi.org/10.1128/AEM.01837-08>
- De Miccolis Angelini RM, Rotolo C, Gerin D, Abate D, Pollastro S, Faretra F (2019) Global transcriptome analysis and differentially expressed genes in grapevine after application of the yeast-derived defense inducer cerevisane. *Pest Manag Sci* 75:2020–2033. <https://doi.org/10.1002/ps.5317>
- d'Enfert C, Janbon G (2016) Biofilm formation in *Candida glabrata*: what have we learnt from functional genomics approaches? *FEMS Yeast Res* 16:fov111. <https://doi.org/10.1093/femsyr/fov111>
- Desai JV, Mitchell AP, Andes DR (2014) Fungal biofilms, drug resistance, and recurrent infection. *Cold Spring Harb Perspect Med*. <https://doi.org/10.1101/cshperspect.a019729>
- Dhami MK, Hartwig T, Fukami T (2016) Genetic basis of priority effects: insights from nectar yeast. *Proc Biol Sci*. <https://doi.org/10.1098/rspb.2016.1455>
- Di Francesco A, Ugolini L, Lazzeri L, Mari M (2014) Production of volatile organic compounds by *Aureobasidium pullulans* as a potential mechanism of action against postharvest fruit pathogens. *Biol Control* 81:8–14. <https://doi.org/10.1016/j.biocontrol.2014.10.004>
- Dogi CA, Armando R, Luduena R, de Moreno de LeBlanc A, Rosa CA, Dalcero A, Cavaglieri L (2011) *Saccharomyces cerevisiae* strains retain their viability and aflatoxin B1 binding ability under gastrointestinal conditions and improve ruminal fermentation. *Food Addit Contam Part A* 28:1705–1711. <https://doi.org/10.1080/19440049.2011.605771>
- Droby S, Chalutz E (1994) Mode of action of biocontrol agents of postharvest diseases. In: Wilson CL, Wisniewski ME (eds) *Biological control of postharvest diseases: theory and practice*. CRC Press Inc, Boca Raton, pp 63–75
- Droby S, El-Gerberia B (2006) Yeast *Metschnikowia fructicola* NRRL Y-30752 for inhibiting deleterious microorganisms on plants. USA Patent, 7 Feb 2006
- Droby S et al (1998) Commercial testing of Aspire: a yeast preparation for the biological control of postharvest decay of citrus. *Biol Control* 12:97–101
- Droby S, Vinokur V, Weiss B, Cohen L, Daus A, Goldschmidt EE, Porat R (2002) Induction of resistance to *Penicillium digitatum* in grapefruit by the yeast biocontrol agent *Candida oleophila*. *Phytopathology* 92:393–399. <https://doi.org/10.1094/PHYTO.2002.92.4.393>
- Droby S, Wisniewski M, El-Ghaouth A, Wilson C (2003) Biological control of postharvest diseases of fruit and vegetables: current

- achievements and future challenges. In: International Society for Horticultural Science (ISHS), Leuven, Belgium, pp 703–713. <https://doi.org/10.17660/actahortic.2003.628.89>
- Droby S, Wisniewski M, El Ghaouth A, Wilson C (2003b) Influence of food additives on the control of postharvest rots of apple and peach and efficacy of the yeast-based biocontrol product aspire. *Postharvest Biol Tec* 27:127–135. [https://doi.org/10.1016/S0925-5214\(02\)00046-7](https://doi.org/10.1016/S0925-5214(02)00046-7)
- Droby S, Wisniewski M, Macarasin D, Wilson C (2009) Twenty years of postharvest biocontrol research: is it time for a new paradigm? *Postharvest Biol Technol* 52:137–145. <https://doi.org/10.1016/j.postharvbio.2008.11.009>
- El Ghaouth A, Wilson CL, Wisniewski M (2003) Control of postharvest decay of apple fruit with *Candida saitoana* and induction of defense responses. *Phytopathology* 93:344–348. <https://doi.org/10.1094/PHYTO.2003.93.3.344>
- Elad Y, Köhl J, Fokkema N (1994) Control of infection and sporulation of *Botrytis cinerea* on bean and tomato by saprophytic yeasts. *Phytopathology* 84:1193–1200
- El-Ghaouth A, Wilson C (2002) *Candida saitoana* compositions for biocontrol of plant postharvest decay USA Patent
- El-Ghaouth A, Wilson CL, Wisniewski M (1998) Ultrastructural and cytochemical aspects of the biological control of *Botrytis cinerea* by *Candida saitoana* in apple fruit. *Phytopathology* 88:282–291. <https://doi.org/10.1094/PHYTO.1998.88.4.282>
- El-Ghaouth A, Smilanick JL, Brown GE, Ippolito A, Wisniewski M, Wilson CL (2000a) Application of *Candida saitoana* and glycolchitosan for the control of postharvest diseases of apple and citrus fruit under semi-commercial conditions. *Plant Dis* 84:243–248. <https://doi.org/10.1094/Pdis.2000.84.3.243>
- El-Ghaouth A, Smilanick JL, Wilson CL (2000b) Enhancement of the performance of *Candida saitoana* by the addition of glycolchitosan for the control of postharvest decay of apple and citrus fruit. *Postharvest Biol Technol* 19:103–110. [https://doi.org/10.1016/S0925-5214\(00\)00076-4](https://doi.org/10.1016/S0925-5214(00)00076-4)
- El-Ghaouth A, Smilanick JL, Wisniewski M, Wilson CL (2000c) Improved control of apple and citrus fruit decay with a combination of *Candida saitoana* and 2-deoxy-D-glucose. *Plant Dis* 84:249–253. <https://doi.org/10.1094/Pdis.2000.84.3.249>
- Eshel D, Regev R, Orenstein J, Droby S, Gan-Mor S (2009) Combining physical, chemical and biological methods for synergistic control of postharvest diseases: a case study of black root rot of carrot. *Postharvest Biol Technol* 54:48–52. <https://doi.org/10.1016/j.postharvbio.2009.04.011>
- European Commission Health & Consumers Directorate-General (2013) Review report for the active substance *Candida oleophila* strain O. vol SANCO/10395/2013 rev 1. <https://doi.org/10.2903/j.efsa.2015.4250>
- European Food Safety Authority (2005) Opinion of the Scientific Committee on a request from EFSA related to a generic approach to the safety assessment by EFSA of microorganisms used in food/feed and the production of food/feed additives. *EFSA J* 3:226. <https://doi.org/10.2903/j.efsa.2005.226>
- European Food Safety Authority (EFSA) (2013) Conclusion on the peer review of the pesticide risk assessment of the active substance *Aureobasidium pullulans* (strains DSM 14940 and DSM 14941). *EFSA J* 11:3183. <https://doi.org/10.2903/j.efsa.2015.4322>
- European Food Safety Authority (EFSA) (2015a) Conclusion on the peer review of the pesticide risk assessment of the active substance *Candida oleophila* strain O. *EFSA J* 10:2944. <https://doi.org/10.2903/j.efsa.2015.4250>
- European Food Safety Authority (EFSA) (2015b) Peer review of the pesticide risk assessment of the active substance *Saccharomyces cerevisiae* strain LAS02. *EFSA J* 13:4322. <https://doi.org/10.2903/j.efsa.2015.4322>
- European Food Safety Authority (EFSA) (2015c) Public consultation on the active substance *Metschnikowia fructicola* NRRL Y-27328 EFSA Rapporteur Assessment Report
- European Food Safety Authority (EFSA) (2017) Peer review of the pesticide risk assessment of the active substance *Metschnikowia fructicola* NRRL Y-27328. *EFSA J* 15:5084. <https://doi.org/10.2903/j.efsa.2017.5084>
- Fan Q, Tian S (2001) Postharvest biological control of grey mold and blue mold on apple by *Cryptococcus albidus* (Saito) Skinner. *Postharvest Biol Technol* 21:341–350. [https://doi.org/10.1016/S0925-5214\(00\)00182-4](https://doi.org/10.1016/S0925-5214(00)00182-4)
- Fanning S, Mitchell AP (2012) Fungal biofilms. *PLoS Pathog* 8:e1002585. <https://doi.org/10.1371/journal.ppat.1002585>
- Farbo MG et al (2018) Effect of yeast volatile organic compounds on ochratoxin A-producing *Aspergillus carbonarius* and *A. ochraceus*. *Int J Food Microbiol* 284:1–10. <https://doi.org/10.1016/j.ijfoodmicro.2018.06.023>
- Fialho MB, Toffano L, Pedroso MP, Augusto F, Pascholati SF (2010) Volatile organic compounds produced by *Saccharomyces cerevisiae* inhibit the *in vitro* development of *Guignardia citricarpa*, the causal agent of citrus black spot. *World J Microb Biot* 26:925–932. <https://doi.org/10.1007/s11274-009-0255-4>
- Fiori S, Fadda A, Giobbe S, Berardi E, Migheli Q (2008) *Pichia angusta* is an effective biocontrol yeast against postharvest decay of apple fruit caused by *Botrytis cinerea* and *Monilia fructicola*. *FEMS Yeast Res* 8:961–963. <https://doi.org/10.1111/j.1567-1364.2008.00424.x>
- Fiori S et al (2012) Identification of differentially expressed genes associated with changes in the morphology of *Pichia fermentans* on apple and peach fruit. *FEMS Yeast Res* 12:785–795. <https://doi.org/10.1111/j.1567-1364.2012.00829.x>
- Fiori S, Urgeghe PP, Hammami W, Razzu S, Jaoua S, Migheli Q (2014) Biocontrol activity of four non- and low-fermenting yeast strains against *Aspergillus carbonarius* and their ability to remove ochratoxin A from grape juice. *Int J Food Microbiol* 189:45–50. <https://doi.org/10.1016/j.ijfoodmicro.2014.07.020>
- Fitzpatrick DA (2012) Horizontal gene transfer in fungi. *FEMS Microbiol Lett* 329:1–8. <https://doi.org/10.1111/j.1574-6968.2011.02465.x>
- Friel D, Pessoa NM, Vandenbol M, Jijakli MH (2007) Separate and combined disruptions of two *exo-β-1,3-glucanase* genes decrease the efficiency of *Pichia anomala* (strain K) biocontrol against *Botrytis cinerea* on apple. *Mol Plant Microbe Interact* 20:371–379. <https://doi.org/10.1094/MPMI-20-4-0371>
- Gafni N, Calderon CE, Harris R, Buxdorf K, Dafa-Berger A, Zeilinger-Reichert E, Levy M (2015) Biological control of the cucurbit powdery mildew pathogen *Podosphaera xanthii* by means of the epiphytic fungus *Pseudozyma aphidis* and parasitism as a mode of action. *Front Plant Sci* 6:132. <https://doi.org/10.3389/fpls.2015.00132>
- Gamagae SU, Sivakumar D, Wijeratnam RSW, Wijesundera RLC (2003) Use of sodium bicarbonate and *Candida oleophila* to control anthracnose in papaya during storage. *Crop Prot* 22:775–779. [https://doi.org/10.1016/S0261-2194\(03\)00046-2](https://doi.org/10.1016/S0261-2194(03)00046-2)
- Gauthier GM (2015) Dimorphism in fungal pathogens of mammals, plants, and insects. *PLoS Pathog* 11:e1004608. <https://doi.org/10.1371/journal.ppat.1004608>
- Gauthier GM (2017) Fungal dimorphism and virulence: molecular mechanisms for temperature adaptation, immune evasion, and *in vivo* survival. *Mediators Inflamm* 2017:8491383. <https://doi.org/10.1155/2017/8491383>
- Giobbe S, Marceddu S, Scherm B, Zara G, Mazzarello VL, Budroni M, Migheli Q (2007) The strange case of a biofilm-forming strain of *Pichia fermentans*, which controls *Monilinia* brown rot on apple but is pathogenic on peach fruit. *FEMS Yeast Res* 7:1389–1398. <https://doi.org/10.1111/j.1567-1364.2007.00301.x>

- Golubev WI, Pfeiffer I, Golubeva EW (2006) Mycocin production in *Pseudozyma tsukubaensis*. *Mycopathologia* 162:313–316. <https://doi.org/10.1007/s11046-006-0065-2>
- Goold HD, Wright P, Hailstones D (2018) Emerging opportunities for synthetic biology in agriculture. *Genes (Basel)* 9:E341. <https://doi.org/10.3390/genes9070341>
- Gore-Lloyd D et al (2019) Snf2 controls pulcherriminic acid biosynthesis and antifungal activity of the biocontrol yeast *Metschnikowia pulcherrima*. *Mol Microbiol* 112:317–332. <https://doi.org/10.1111/mmi.14272>
- Gozzo F, Faoro F (2013) Systemic acquired resistance (50 years after discovery): moving from the lab to the field. *J Agric Food Chem* 61:12473–12491. <https://doi.org/10.1021/jf401456x>
- Grevesse C, Lepoivre P, Jijakli MH (2003) Characterization of the exoglucanase-encoding gene *PaEXG2* and study of its role in the biocontrol activity of *Pichia anomala* strain K. *Phytopathology* 93:1145–1152. <https://doi.org/10.1094/PHYTO.2003.93.9.1145>
- Guo FJ, Ma Y, Xu HM, Wang XH, Chi ZM (2013) A novel killer toxin produced by the marine-derived yeast *Wickerhamomyces anomalus* YF07b. *Antonie Van Leeuwenhoek* 103:737–746. <https://doi.org/10.1007/s10482-012-9855-3>
- Guyard C, Dehecq E, Tissier JP, Polonelli L, Dei-Cas E, Cailliez JC, Menozzi FD (2002a) Involvement of [beta]-glucans in the wide-spectrum antimicrobial activity of *Williopsis saturnus* var. *mrakii* MUCL 41968 killer toxin. *Mol Med* 8:686–694
- Guyard C et al (2002b) Characterization of a *Williopsis saturnus* var. *mrakii* high molecular weight secreted killer toxin with broad-spectrum antimicrobial activity. *J Antimicrob Chemother* 49:961–971. <https://doi.org/10.1093/jac/dkf040>
- Hadwiger LA, McDonel H, Glawe D (2015) Wild yeast strains as prospective candidates to induce resistance against potato late blight (*Phytophthora infestans*). *Am J Potato Res* 92:379–386. <https://doi.org/10.1007/s12230-015-9443-y>
- Helbig J (2002) Ability of the antagonistic yeast *Cryptococcus albidus* to control *Botrytis cinerea* in strawberry. *Biocontrol* 47:85–99. <https://doi.org/10.1023/A:1014466903941>
- Hernandez A, Martin A, Aranda E, Perez-Navado F, Cordoba MG (2007) Identification and characterization of yeast isolated from the elaboration of seasoned green table olives. *Food Microbiol* 24:346–351. <https://doi.org/10.1016/j.fm.2006.07.022>
- Herrera-Estrella A, Chet I (1999) Chitinases in biological control. *EXS* 87:171–184
- Hershkovitz V et al (2012) Global changes in gene expression of grapefruit peel tissue in response to the yeast biocontrol agent *Metschnikowia fructicola*. *Mol Plant Pathol* 13:338–349. <https://doi.org/10.1111/j.1364-3703.2011.00750.x>
- Hershkovitz V et al (2013) De-novo assembly and characterization of the transcriptome of *Metschnikowia fructicola* reveals differences in gene expression following interaction with *Penicillium digitatum* and grapefruit peel. *BMC Genom* 14:168. <https://doi.org/10.1186/1471-2164-14-168>
- Heydari A, Pessarakli M (2010) A review on biological control of fungal plant pathogens using microbial agents. *J Biol Sci* 10:273–290. <https://doi.org/10.3923/jbs.2010.273.290>
- Hilber-Bodmer M, Schmid M, Ahrens CH, Freimoser FM (2017) Competition assays and physiological experiments of soil and phyllosphere yeasts identify *Candida subhashii* as a novel antagonist of filamentous fungi. *BMC Microbiol* 17:4. <https://doi.org/10.1186/s12866-016-0908-z>
- Holb IJ, Kunz S (2013) Integrated control of brown rot blossom blight by combining approved chemical control options with *Aureobasidium pullulans* in organic cherry production. *Crop Prot* 54:114–120. <https://doi.org/10.1016/j.cropro.2013.07.003>
- Hua MX, Chi Z, Liu GL, Buzdar MA, Chi ZM (2010) Production of a novel and cold-active killer toxin by *Mrakia frigida* 2E00797 isolated from sea sediment in Antarctica. *Extremophiles* 14:515–521. <https://doi.org/10.1007/s00792-010-0331-6>
- Hua SS, Beck JJ, Sarreal SB, Gee W (2014) The major volatile compound 2-phenylethanol from the biocontrol yeast, *Pichia anomala*, inhibits growth and expression of aflatoxin biosynthetic genes of *Aspergillus flavus*. *Mycotoxin Res* 30:71–78. <https://doi.org/10.1007/s12550-014-0189-z>
- Huang R, Li GQ, Zhang J, Yang L, Che HJ, Jiang DH, Huang HC (2011) Control of postharvest *Botrytis* fruit rot of strawberry by volatile organic compounds of *Candida intermedia*. *Phytopathology* 101:859–869. <https://doi.org/10.1094/PHYTO-09-10-0255>
- Huang R, Che HJ, Zhang J, Yang L, Jiang DH, Li GQ (2012) Evaluation of *Sporidiobolus pararoseus* strain YCXT3 as biocontrol agent of *Botrytis cinerea* on post-harvest strawberry fruits. *Biol Control* 62:53–63. <https://doi.org/10.1016/j.biocontrol.2012.02.010>
- Janisiewicz WJ, Pereira IB, Almeida MS, Roberts DP, Wisniewski M, Kurtenbach E (2008) Improved biocontrol of fruit decay fungi with *Pichia pastoris* recombinant strains expressing Psd1 antifungal peptide. *Postharvest Biol Technol* 47:218–225. <https://doi.org/10.1016/j.postharvbio.2007.06.010>
- Jiang B, Ram AF, Sheraton J, Klis FM, Bussey H (1995) Regulation of cell wall beta-glucan assembly: *pTC1* negatively affects PBS2 action in a pathway that includes modulation of *EXG1* transcription. *Mol Gen Genet* 248:260–269
- Johnson L (2008) Iron and siderophores in fungal-host interactions. *Mycol Res* 112:170–183. <https://doi.org/10.1016/j.mycres.2007.11.012>
- Jones JD, Dangl JL (2006) The plant immune system. *Nature* 444:323–329. <https://doi.org/10.1038/nature05286>
- Jones RW, Prusky D (2002) Expression of an antifungal peptide in *Saccharomyces*: a new approach for biological control of the postharvest disease caused by *Colletotrichum coccodes*. *Phytopathology* 92:33–37. <https://doi.org/10.1094/PHYTO.2002.92.1.33>
- Junker K, Hesselbart A, Wendland J (2017) Draft genome sequence of *Saccharomycopsis fodiens* CBS 8332, a necrotrophic mycoparasite with biocontrol potential. *Genome Announc*. <https://doi.org/10.1128/genomeA.01278-17>
- Junker K, Bravo Ruiz G, Lorenz A, Walker L, Gow NAR, Wendland J (2018) The mycoparasitic yeast *Saccharomycopsis schoenii* predates and kills multi-drug resistant *Candida auris*. *Sci Rep* 8:14959. <https://doi.org/10.1038/s41598-018-33199-z>
- Junker K, Chailyan A, Hesselbart A, Forster J, Wendland J (2019) Multi-omics characterization of the necrotrophic mycoparasite *Saccharomycopsis schoenii*. *PLoS Pathog* 15:e1007692. <https://doi.org/10.1371/journal.ppat.1007692>
- Kasahara S et al (1994a) Involvement of cell wall beta-glucan in the action of HM-1 killer toxin. *FEBS Lett* 348:27–32
- Kasahara S et al (1994b) Cloning of the *Saccharomyces cerevisiae* gene whose overexpression overcomes the effects of HM-1 killer toxin, which inhibits beta-glucan synthesis. *J Bacteriol* 176:1488–1499. <https://doi.org/10.1128/jb.176.5.1488-1499.1994>
- Keyhani NO (2018) Lipid biology in fungal stress and virulence: entomopathogenic fungi. *Fungal Biol* 122:420–429. <https://doi.org/10.1016/j.funbio.2017.07.003>
- Klassen R, Teichert S, Meinhardt F (2004) Novel yeast killer toxins provoke S-phase arrest and DNA damage checkpoint activation. *Mol Microbiol* 53:263–273. <https://doi.org/10.1111/j.1365-2958.2004.04119.x>
- Klein MN, Kupper KC (2018) Biofilm production by *Aureobasidium pullulans* improves biocontrol against sour rot in citrus. *Food Microbiol* 69:1–10. <https://doi.org/10.1016/j.fm.2017.07.008>
- Klemsdal SS, Tronsmo A (1999) Genetic manipulation for improvement of microbial biocontrol agents. In: Albajes R, Gullino ML, van Lenteren JC, Elad Y (eds) *Integrated pest and disease*

- management in greenhouse crops. Springer, Dordrecht, pp 353–364. https://doi.org/10.1007/0-306-47585-5_25
- Kombrink A, Sanchez-Vallet A, Thomma BP (2011) The role of chitin detection in plant–pathogen interactions. *Microbes Infect* 13:1168–1176. <https://doi.org/10.1016/j.micinf.2011.07.010>
- Kowalska J, Drożdżyński D, Remlein-Starosta D, Sas-Paszcz L, Malusá E (2012) Use of *Cryptococcus albidus* for controlling grey mould in the production and storage of organically grown strawberries. *J Plant Dis Protect* 119:174–178
- Kunz S (2004) Development of “Blossom-Protect” - a yeast preparation for the reduction of blossom infections by fire blight. 11th International Conference on Cultivation Technique and Phytopathological problems in organic fruit-growing, Weinsberg, Germany
- Kunz S, Haug P (2006) Development of a strategy for fire blight control in organic fruit growing. In: Boos M (ed) 12th International Conference on Cultivation Technique and Phytopathological Problems in Organic Fruit-Growing, Weinsberg, 2006. Fördergemeinschaft ökologischer Obstbau, pp 113–117
- Kunz S, Schmitt A, Haug P (2011) Field testing of strategies for fire blight control in organic fruit growing. *Acta Hort* 896:431–436. <https://doi.org/10.17660/ActaHort.2011.896.62>
- Kurtzman CP, Droby S (2001) *Metschnikowia fructicola*, a new ascospore yeast with potential for biocontrol of postharvest fruit rots. *Syst Appl Microbiol* 24:395–399. <https://doi.org/10.1078/0723-2020-00045>
- Lachance MA, Pang WM (1997) Predacious yeasts. *Yeast* 13:225–232. [https://doi.org/10.1002/\(SICI\)1097-0061\(19970315\)13:3%3c225::AID-YEA87%3e3.0.CO;2-I](https://doi.org/10.1002/(SICI)1097-0061(19970315)13:3%3c225::AID-YEA87%3e3.0.CO;2-I)
- Lachance MA, Starmer WT, Rosa CA, Bowles JM, Barker JS, Janzen DH (2001) Biogeography of the yeasts of ephemeral flowers and their insects. *FEMS Yeast Res* 1:1–8. <https://doi.org/10.1111/j.1567-1364.2001.tb00007.x>
- Lahlali R, Serrhini MN, Jijakli MH (2004) Efficacy assessment of *Candida oleophila* (strain O) and *Pichia anomala* (strain K) against major postharvest diseases of citrus fruits in Morocco. *Commun Agric Appl Biol Sci* 69:601–609
- Langner T, Gohre V (2015) Fungal chitinases: function, regulation, and potential roles in plant/pathogen interactions. *Curr Genet*. <https://doi.org/10.1007/s00294-015-0530-x>
- Lee G, Lee SH, Kim KM, Ryu CM (2017) Foliar application of the leaf-colonizing yeast *Pseudozyma churashimaensis* elicits systemic defense of pepper against bacterial and viral pathogens. *Sci Rep* 7:39432. <https://doi.org/10.1038/srep39432>
- Lemos WJ Jr et al (2016) Biocontrol ability and action mechanism of *Starmarella bacillaris* (synonym *Candida zemlinina*) isolated from wine musts against gray mold disease agent *Botrytis cinerea* on grape and their effects on alcoholic fermentation. *Front Microbiol* 7:1249. <https://doi.org/10.3389/fmicb.2016.01249>
- Li G et al (2016) Stress tolerance and biocontrol performance of the yeast antagonist, *Candida diversa*, change with morphology transition. *Environ Sci Pollut Res Int* 23:2962–2967. <https://doi.org/10.1007/s11356-015-5769-8>
- Liu HM, Guo JH, Liu P, Cheng YJ, Wang BQ, Long CA, Deng BX (2010) Inhibitory activity of tea polyphenol and *Candida ernobii* against *Diplodia natalensis* infections. *J Appl Microbiol* 108:1066–1072. <https://doi.org/10.1111/j.1365-2672.2009.04511.x>
- Liu T et al (2012) Chitin-induced dimerization activates a plant immune receptor. *Science* 336:1160–1164. <https://doi.org/10.1126/science.1218867>
- Liu X et al (2014) Host-induced bacterial cell wall decomposition mediates pattern-triggered immunity in *Arabidopsis*. *Elife*. <https://doi.org/10.7554/eLife.01990>
- Liu GL, Chi Z, Wang GY, Wang ZP, Li Y, Chi ZM (2015) Yeast killer toxins, molecular mechanisms of their action and their applications. *Crit Rev Biotechnol* 35:222–234. <https://doi.org/10.3109/07388551.2013.833582>
- Liu P, Chen K, Li G, Yang X, Long CA (2016) Comparative transcriptional profiling of orange fruit in response to the biocontrol yeast *Kloeckera apiculata* and its active compounds. *BMC Genom* 17:17. <https://doi.org/10.1186/s12864-015-2333-3>
- Liu J, Li G, Sui Y (2017) Optimization of culture medium enhances viable biomass production and biocontrol efficacy of the antagonistic yeast, *Candida diversa*. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2017.02021>
- Lohse MB, Gulati M, Johnson AD, Nobile CJ (2018) Development and regulation of single- and multi-species *Candida albicans* biofilms. *Nat Rev Microbiol* 16:19–31. <https://doi.org/10.1038/nrmicro.2017.107>
- Lopes FAC et al (2012) Biochemical and metabolic profiles of *Trichoderma* strains isolated from common bean crops in the Brazilian Cerrado, and potential antagonism against *Sclerotinia sclerotiorum*. *Fungal Biol* 116:815–824. <https://doi.org/10.1016/j.funbio.2012.04.015>
- Lopes MR, Klein MN, Ferraz LP, da Silva AC, Kupper KC (2015) *Saccharomyces cerevisiae*: a novel and efficient biological control agent for *Colletotrichum acutatum* during pre-harvest. *Microbiol Res* 175:93–99. <https://doi.org/10.1016/j.micres.2015.04.003>
- Lowes KF, Shearman CA, Payne J, MacKenzie D, Archer DB, Merry RJ, Gasson MJ (2000) Prevention of yeast spoilage in feed and food by the yeast mycocin HMK. *Appl Environ Microbiol* 66:1066–1076
- Lu L et al (2013) Preharvest application of antagonistic yeast *Rhodospodium paludigenum* induced resistance against postharvest diseases in mandarin orange. *Biol Control* 67:130–136. <https://doi.org/10.1016/j.biocontrol.2013.07.016>
- Lu L, Xu S, Zeng L, Zheng X, Yu T (2014) *Rhodospodium paludigenum* induced resistance in Ponkan mandarin against *Penicillium digitatum* requires ethylene-dependent signaling pathway. *Postharvest Biol Technol* 97:93–101. <https://doi.org/10.1016/j.postharvbio.2014.06.007>
- Luksa J, Podoliankaite M, Vepstaite I, Strazdaite-Zieliene Z, Urbonavičius J, Serviėne E (2015) Yeast beta-1,6-glucan is a primary target for the *Saccharomyces cerevisiae* K2 toxin. *Eukaryot Cell* 14:406–414. <https://doi.org/10.1128/EC.00287-14>
- Lutz MC, Lopes CA, Rodriguez ME, Sosa MC, Sangorrin MP (2013) Efficacy and putative mode of action of native and commercial antagonistic yeasts against postharvest pathogens of pear. *Int J Food Microbiol* 164:166–172. <https://doi.org/10.1016/j.ijfoodmicro.2013.04.005>
- Macarisin D, Droby S, Bauchan G, Wisniewski M (2010) Superoxide anion and hydrogen peroxide in the yeast antagonist-fruit interaction: a new role for reactive oxygen species in postharvest biocontrol? *Postharvest Biol Technol* 58:194–202. <https://doi.org/10.1016/j.postharvbio.2010.07.008>
- Mannazzu I et al (2019) Yeast killer toxins: from ecological significance to application. *Crit Rev Biotechnol* 39:603–617. <https://doi.org/10.1080/07388551.2019.1601679>
- Manuel J, Selin C, Fernando WG, de Kievit T (2012) Stringent response mutants of *Pseudomonas chlororaphis* PA23 exhibit enhanced antifungal activity against *Sclerotinia sclerotiorum* in vitro. *Microbiology* 158:207–216. <https://doi.org/10.1099/mic.0.053082-0>
- Marchand G, Clement-Mathieu G, Neveu B, Belanger RR (2007) Enhancing biological control efficacy of yeasts to control fungal diseases through biotechnology. In: Punja ZK, De Boer SH, Sanfaçon H (eds) *Biotechnology and plant disease management*. CAB International, Wallingford, pp 518–531. <https://doi.org/10.1079/9781845932886.0518>

- Marquina D, Barroso J, Santos A, Peinado JM (2001) Production and characteristics of *Debaryomyces hansenii* killer toxin. *Microbiol Res* 156:387–391. <https://doi.org/10.1078/0944-5013-00117>
- Marquina D, Santos A, Peinado JM (2002) Biology of killer yeasts. *Int Microbiol* 5:65–71. <https://doi.org/10.1007/s10123-002-0066-z>
- Maselko M, Heinsch SC, Chacon JM, Harcombe WR, Smanski MJ (2017) Engineering species-like barriers to sexual reproduction. *Nat Commun* 8:883. <https://doi.org/10.1038/s41467-017-01007-3>
- Maserti B, Podda A, Giorgetti L, Del Carratore R, Chevret D, Migheli Q (2015) Proteome changes during yeast-like and pseudohyphal growth in the biofilm-forming yeast *Pichia fermentans*. *Amino Acids* 47:1091–1106. <https://doi.org/10.1007/s0072-6-015-1933-1>
- Masoud W, Poll L, Jakobsen M (2005) Influence of volatile compounds produced by yeasts predominant during processing of *Coffea arabica* in East Africa on growth and ochratoxin A (OTA) production by *Aspergillus ochraceus*. *Yeast* 22:1133–1142. <https://doi.org/10.1002/yea.1304>
- Mayer FL, Wilson D, Hube B (2013) *Candida albicans* pathogenicity mechanisms. *Virulence* 4:119–128. <https://doi.org/10.4161/viru.22913>
- Mbili NC (2012) Evaluation of integrated control of postharvest grey mould and blue mould of pome fruit using yeast, potassium silicate and hot water treatments. University of KwaZulu-Natal
- McCormack P, Wildman HG, Jeffries P (1995) The influence of moisture on the suppression of *Pseudomonas syringae* by *Aureobasidium pullulans* on an artificial leaf surface. *FEMS Microbiol Ecol* 16:159–165. [https://doi.org/10.1016/0168-6496\(95\)92762-Z](https://doi.org/10.1016/0168-6496(95)92762-Z)
- McLaughlin RJ, Wilson CL, Droby S, Benarie R, Chalutz E (1992) Biological control of postharvest diseases of grape, peach, and apple with the yeasts *Kloeckera apiculata* and *Candida guilliermondii*. *Plant Dis* 76:470–473
- Miceli MH, Diaz JA, Lee SA (2011) Emerging opportunistic yeast infections. *Lancet Infect Dis* 11:142–151. [https://doi.org/10.1016/S1473-3099\(10\)70218-8](https://doi.org/10.1016/S1473-3099(10)70218-8)
- Mimee B, Labbe C, Pelletier R, Belanger RR (2005) Antifungal activity of flocculosin, a novel glycolipid isolated from *Pseudozyma flocculosa*. *Antimicrob Agents Chemother* 49:1597–1599. <https://doi.org/10.1128/AAC.49.4.1597-1599.2005>
- Mimee B, Labbe C, Belanger RR (2009) Catabolism of flocculosin, an antimicrobial metabolite produced by *Pseudozyma flocculosa*. *Glycobiology* 19:995–1001. <https://doi.org/10.1093/glyco/b/cwp078>
- Molinu MG, Pani G, Venditti T, Dore A, Ladu G, D'Hallewin G (2011) Sequential application of NaHCO₃, CaCl₂ and *Candida oleophila* (isolate 13L) affects significantly *Penicillium expansum* growth and the infection degree in apples. *Commun Agric Appl Biol Sci* 76:743–750
- Morales H, Sanchis V, Usall J, Ramos AJ, Marin S (2008) Effect of biocontrol agents *Candida sake* and *Pantoea agglomerans* on *Penicillium expansum* growth and patulin accumulation in apples. *Int J Food Microbiol* 122:61–67. <https://doi.org/10.1016/j.ijfoodmicro.2007.11.056>
- Morath SU, Hung R, Bennett JW (2012) Fungal volatile organic compounds: a review with emphasis on their biotechnological potential. *Fungal Biol Rev* 26:73–83. <https://doi.org/10.1016/j.fbr.2012.07.001>
- Moriguchi K, Yamamoto S, Tanaka K, Kurata N, Suzuki K (2013) Trans-kingdom horizontal DNA transfer from bacteria to yeast is highly plastic due to natural polymorphisms in auxiliary nonessential recipient genes. *PLoS ONE* 8:e74590. <https://doi.org/10.1371/journal.pone.0074590>
- Nagpure A, Choudhary B, Gupta RK (2014) Chitinases: in agriculture and human healthcare. *Crit Rev Biotechnol* 34:215–232. <https://doi.org/10.3109/07388551.2013.790874>
- Nigro F, Finetti Sialer MM, Gallitelli D (1999) Transformation of *Metschnikowia pulcherrima* 320, biocontrol agent of storage rot, with the green fluorescent protein gene. *J Plant Pathol* 81:205–208
- Nunes C, Usall J, Teixido N, Abadias M, Viñas I (2002a) Improved control of postharvest decay of pears by the combination of *Candida sake* (CPA-1) and ammonium molybdate. *Phytopathology* 92:281–287. <https://doi.org/10.1094/Phyto.2002.92.3.281>
- Nunes C, Usall J, Teixido N, Viñas I (2002b) Improvement of *Candida sake* biocontrol activity against post-harvest decay by the addition of ammonium molybdate. *J Appl Microbiol* 92:927–935
- Oporto CI, Villarroel CA, Tapia SM, Garcia V, Cubillos FA (2019) Distinct transcriptional changes in response to patulin underlie toxin biosorption differences in *Saccharomyces cerevisiae*. *Toxins (Basel)*. <https://doi.org/10.3390/toxins11070400>
- Opulente DA et al (2019) Pathogenic budding yeasts isolated outside of clinical settings. *FEMS Yeast Res*. <https://doi.org/10.1093/femsyr/foz032>
- Oro L, Feliziani E, Ciani M, Romanazzi G, Comitini F (2017) Volatile organic compounds from *Wickerhamomyces anomalus*, *Metschnikowia pulcherrima* and *Saccharomyces cerevisiae* inhibit growth of decay causing fungi and control postharvest diseases of strawberries. *Int J Food Microbiol* 265:18–22. <https://doi.org/10.1016/j.ijfoodmicro.2017.10.027>
- Ortu G, Demontis MA, Budroni M, Goyard S, d'Enfert C, Migheli Q (2005) Study of biofilm formation in *Candida albicans* may help understanding the biocontrol capability of a *flor* strain of *Saccharomyces cerevisiae* against the phytopathogenic fungus *Penicillium expansum*. *J Plant Pathol* 87:300
- Pandin C, Le Coq D, Canette A, Aymerich S, Briandet R (2017) Should the biofilm mode of life be taken into consideration for microbial biocontrol agents? *Microb Biotechnol* 10:719–734. <https://doi.org/10.1111/1751-7915.12693>
- Papanikolaou S, Aggelis G (2011) Lipids of oleaginous yeasts. Part I: biochemistry of single cell oil production. *Eur J Lipid Sci Technol* 113:1031–1051. <https://doi.org/10.1002/ejlt.201100014>
- Papon N et al (2013) *Candida guilliermondii*: biotechnological applications, perspectives for biological control, emerging clinical importance and recent advances in genetics. *Curr Genet* 59:73–90. <https://doi.org/10.1007/s00294-013-0391-0>
- Parafati L, Vitale A, Restuccia C, Cirvilleri G (2015) Biocontrol ability and action mechanism of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape. *Food Microbiol* 47:85–92. <https://doi.org/10.1016/j.fm.2014.11.013>
- Parafati L, Cirvilleri G, Restuccia C, Wisniewski M (2017a) Potential role of exoglucanase genes (*WaEXG1* and *WaEXG2*) in the biocontrol activity of *Wickerhamomyces anomalus*. *Microb Ecol* 73:876–884. <https://doi.org/10.1007/s00248-016-0887-5>
- Parafati L, Vitale A, Restuccia C, Cirvilleri G (2017b) Performance evaluation of volatile organic compounds by antagonistic yeasts immobilized on hydrogel spheres against gray, green and blue postharvest decays. *Food Microbiol* 63:191–198. <https://doi.org/10.1016/j.fm.2016.11.021>
- Park M, Do E, Jung WH (2013) Lipolytic enzymes involved in the virulence of human pathogenic fungi. *Mycobiology* 41:67–72. <https://doi.org/10.5941/MYCO.2013.41.2.67>
- Perez MF et al (2016) Native killer yeasts as biocontrol agents of post-harvest fungal diseases in lemons. *PLoS ONE* 11:e0165590. <https://doi.org/10.1371/journal.pone.0165590>
- Peter J et al (2018) Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature* 556:339–344. <https://doi.org/10.1038/s41586-018-0030-5>
- Piano S, Neyrotti V, Migheli Q, Gullino ML (1997) Biocontrol capability of *Metschnikowia pulcherrima* against *Botrytis* postharvest rot of apple. *Postharvest Biol Technol* 11:131–140. [https://doi.org/10.1016/S0925-5214\(97\)00022-7](https://doi.org/10.1016/S0925-5214(97)00022-7)

- Pieterse CM, Zamioudis C, Berendsen RL, Weller DM, Van Wees SC, Bakker PA (2014) Induced systemic resistance by beneficial microbes. *Annu Rev Phytopathol* 52:347–375. <https://doi.org/10.1146/annurev-phyto-082712-102340>
- Pimenta RS, Silva FL, Silva JF, Morais PB, Braga DT, Rosa CA, Correa A Jr (2008) Biological control of *Penicillium italicum*, *P. digitatum* and *P. expansum* by the predacious yeast *Saccharomyces schoenii* on oranges. *Braz J Microbiol* 39:85–90. <https://doi.org/10.1590/S1517-838220080001000020>
- Piombo E et al (2018) Genome sequence, assembly and characterization of two *Metschnikowia fructicola* strains used as biocontrol agents of postharvest diseases. *Front Microbiol* 9:593. <https://doi.org/10.3389/fmicb.2018.00593>
- Pizzolitto RP, Armando MR, Combina M, Cavagliero LR, Dalcero AM, Salvano MA (2012) Evaluation of *Saccharomyces cerevisiae* strains as probiotic agent with aflatoxin B(1) adsorption ability for use in poultry feedstuffs. *J Environ Sci Health B* 47:933–941. <https://doi.org/10.1080/03601234.2012.706558>
- Pozo MI, Lachance MA, Herrera CM (2012) Nectar yeasts of two southern Spanish plants: the roles of immigration and physiological traits in community assembly. *FEMS Microbiol Ecol* 80:281–293. <https://doi.org/10.1111/j.1574-6941.2011.01286.x>
- Prado G et al (2011) Reduction of aflatoxin B1 in stored peanuts (*Arachis hypogaea* L.) using *Saccharomyces cerevisiae*. *J Food Protect* 74:1003–1006. <https://doi.org/10.4315/0362-028X.JFP-10-380>
- Prasongsuk S, Lotrakul P, Ali I, Bankeeree W, Punnapayak H (2018) The current status of *Aureobasidium pullulans* in biotechnology. *Folia Microbiol (Praha)* 63:129–140. <https://doi.org/10.1007/s12223-017-0561-4>
- Pretschner J et al (2018) Yeasts from different habitats and their potential as biocontrol agents. *Fermentation* 4:31. <https://doi.org/10.3390/fermentation4020031>
- Price NP, Manitchotpisit P, Vermillion KE, Bowman MJ, Leathers TD (2013) Structural characterization of novel extracellular liamocins (mannitol oils) produced by *Aureobasidium pullulans* strain NRRL 50380. *Carbohydr Res* 370:24–32. <https://doi.org/10.1016/j.carres.2013.01.014>
- Price NP, Bischoff KM, Leathers TD, Cosse AA, Manitchotpisit P (2017) Polyols, not sugars, determine the structural diversity of anti-streptococcal liamocins produced by *Aureobasidium pullulans* strain NRRL 50380. *J Antibiot (Tokyo)* 70:136–141. <https://doi.org/10.1038/ja.2016.92>
- Pu L, Jingfan F, Kai C, Chao-an L, Yunjiang C (2014) Phenylethanol promotes adhesion and biofilm formation of the antagonistic yeast *Kloeckera apiculata* for the control of blue mold on citrus. *FEMS Yeast Res* 14:536–546. <https://doi.org/10.1111/1567-1364.12139>
- Punja ZK, Utkhede RS (2003) Using fungi and yeasts to manage vegetable crop diseases. *Trends Biotechnol* 21:400–407. [https://doi.org/10.1016/S0167-7799\(03\)00193-8](https://doi.org/10.1016/S0167-7799(03)00193-8)
- Querol A, Fleet GH (eds) (2006) Yeasts in food and beverages. The Yeast Handbook, vol 2. Springer, Heidelberg
- Ramirez M, Velazquez R, Maqueda M, Lopez-Pineiro A, Ribas JC (2015) A new wine *Torulaspora delbrueckii* killer strain with broad antifungal activity and its toxin-encoding double-stranded RNA virus. *Front Microbiol* 6:983. <https://doi.org/10.3389/fmicb.2015.00983>
- Ren X, Kong Q, Wang H, Yu T, Tang YJ, Zhou WW, Zheng X (2012) Control of apple blue mold by *Pichia pastoris* recombinant strains expressing cecropin A. *Bioprocess Biosyst Eng* 35:761–767. <https://doi.org/10.1007/s00449-011-0656-2>
- Reynolds TB, Fink GR (2001) Bakers' yeast, a model for fungal biofilm formation. *Science* 291:878–881. <https://doi.org/10.1126/science.291.5505.878>
- Richards TA, Leonard G, Soanes D, Talbot N (2011) Gene transfer into the fungi. *Fungal Biol Rev* 25:98–110. <https://doi.org/10.1016/j.fbr.2011.04.003>
- Robert V, Cardinali G, Casadevall A (2015) Distribution and impact of yeast thermal tolerance permissive for mammalian infection. *BMC Biol* 13:18. <https://doi.org/10.1186/s12915-015-0127-3>
- Rodriguez-Cousino N, Maqueda M, Ambrona J, Zamora E, Esteban R, Ramirez M (2011) A new wine *Saccharomyces cerevisiae* killer toxin (Klus), encoded by a double-stranded rna virus, with broad antifungal activity is evolutionarily related to a chromosomal host gene. *Appl Environ Microbiol* 77:1822–1832. <https://doi.org/10.1128/AEM.02501-10>
- Rossouw D, Meiring SP, Bauer FF (2018) Modifying *Saccharomyces cerevisiae* adhesion properties regulates yeast ecosystem dynamics. *mSphere*. <https://doi.org/10.1128/msphere.00383-18>
- Saligkarias ID, Gravanis FT, Epton HAS (2002) Biological control of *Botrytis cinerea* on tomato plants by the use of epiphytic yeasts *Candida guilliermondii* strains 101 and US 7 and *Candida oleophila* strain I-182: I. *in vivo* studies. *Biol Control* 25:143–150. [https://doi.org/10.1016/S1049-9644\(02\)00051-8](https://doi.org/10.1016/S1049-9644(02)00051-8)
- Sánchez-Pérez LdC, Barranco-Florida JE, Rodríguez-Navarro S, Cervantes-Mayagoitia JF, Ramos-Lopez MÁ (2014) Enzymes of entomopathogenic fungi, advances and insights. *Adv Enzyme Res* 2:65–76. <https://doi.org/10.4236/aer.2014.22007>
- Sanna ML, Zara S, Zara G, Migheli Q, Budroni M, Mannazzu I (2012) *Pichia fermentans* dimorphic changes depend on the nitrogen source. *Fungal Biol* 116:769–777. <https://doi.org/10.1016/j.funbio.2012.04.008>
- Sanna ML, Zara G, Zara S, Migheli Q, Budroni M, Mannazzu I (2013) A putative phospholipase C is involved in *Pichia fermentans* dimorphic transition. *Biochim Biophys Acta* 1840:344–349. <https://doi.org/10.1016/j.bbagen.2013.09.030>
- Santos A, Marquina D (2004a) Ion channel activity by *Pichia membranifaciens* killer toxin. *Yeast* 21:151–162. <https://doi.org/10.1002/yea.1069>
- Santos A, Marquina D (2004b) Killer toxin of *Pichia membranifaciens* and its possible use as a biocontrol agent against grey mould disease of grapevine. *Microbiology* 150:2527–2534. <https://doi.org/10.1099/mic.0.27071-0>
- Santos A, Marquina D, Barroso J, Peinado JM (2002) (1→6)-Beta-D-glucan as the cell wall binding site for *Debaryomyces hansenii* killer toxin. *Lett Appl Microbiol* 34:95–99. <https://doi.org/10.1046/j.1472-765x.2002.01053.x>
- Santos A, San Mauro M, Bravo E, Marquina D (2009) PMKT2, a new killer toxin from *Pichia membranifaciens*, and its promising biotechnological properties for control of the spoilage yeast *Brettanomyces bruxellensis*. *Microbiology* 155:624–634. <https://doi.org/10.1099/mic.0.023663-0>
- Saravanakumar D, Clavorella A, Spadaro D, Garibaldi A, Gullino ML (2008) *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and *Penicillium expansum* in apples through iron depletion. *Postharvest Biol Technol* 49:121–128. <https://doi.org/10.1016/j.postharvbio.2007.11.006>
- Saravanakumar D, Spadaro D, Garibaldi A, Gullino ML (2009) Detection of enzymatic activity and partial sequence of a chitinase gene in *Metschnikowia pulcherrima* strain MACH1 used as postharvest biocontrol agent. *Eur J Plant Pathol* 123:183–193. <https://doi.org/10.1007/s10658-008-9355-5>
- Schaible UE, Kaufmann SH (2005) A nutritive view on the host-pathogen interplay. *Trends Microbiol* 13:373–380. <https://doi.org/10.1016/j.tim.2005.06.009>
- Scherm B, Ortu G, Muzzu A, Budroni M, Arras G, Migheli Q (2003) Biocontrol activity of antagonistic yeasts against *Penicillium expansum* on apple. *J Plant Pathol* 85:205–213

- Schmitt MJ, Breinig F (2002) The viral killer system in yeast: from molecular biology to application. *FEMS Microbiol Rev* 26:257–276
- Schmitt MJ, Breinig F (2006) Yeast viral killer toxins: lethality and self-protection. *Nat Rev Microbiol* 4:212–221. <https://doi.org/10.1038/nrmicro1347>
- Schmitt MJ, Schernikau G (1997) Construction of a cDNA-based K1/K2/K28 triple killer strain of *Saccharomyces cerevisiae*. *Food Technol Biotechnol* 35:281–285
- Segal E, Yehuda H, Droby S, Wisniewski M, Goldway M (2002) Cloning and analysis of CoEXG1, a secreted 1,3- β -glucanase of the yeast biocontrol agent *Candida oleophila*. *Yeast* 19:1171–1182. <https://doi.org/10.1002/yea.910>
- Seibold A, Fried A, Kunz S, Moltmann E, Lange E, Jelkmann W (2004) Yeasts as antagonists against fireblight. *EPPO Bull.* <https://doi.org/10.1111/j.1365-2338.2004.00766.x>
- Shalaby ME-S, El-Nady MF (2008) Application of *Saccharomyces cerevisiae* as a biocontrol agent against *Fusarium* infection of sugar beet plants. *Acta Biol Szegediensis* 52:271–275
- Sipiczki M (2006) *Metschnikowia* strains isolated from botrytized grapes antagonize fungal and bacterial growth by iron depletion. *Appl Environ Microb* 72:6716–6724. <https://doi.org/10.1128/Aem.01275-06>
- Slavikova E, Vadkertiova R, Vranova D (2007) Yeasts colonizing the leaf surfaces. *J Basic Microbiol* 47:344–350. <https://doi.org/10.1002/jobm.200710310>
- Sommer B, Overy DP, Haltli B, Kerr RG (2016) Secreted lipases from *Malassezia globosa*: recombinant expression and determination of their substrate specificities. *Microbiology* 162:1069–1079. <https://doi.org/10.1099/mic.0.000299>
- Spadaro D, Droby S (2016) Development of biocontrol products for postharvest diseases of fruit: the importance of elucidating the mechanisms of action of yeast antagonists. *Trends Food Sci Technol* 47:39–49. <https://doi.org/10.1016/j.tifs.2015.11.003>
- Spadaro D, Gullino ML (2005) Improving the efficacy of biocontrol agents against soilborne pathogens. *Crop Prot* 24:601–613. <https://doi.org/10.1016/j.cropro.2004.11.003>
- Spadaro D, Ciavarella A, Dianpeng Z, Garibaldi A, Gullino ML (2010a) Effect of culture media and pH on the biomass production and biocontrol efficacy of a *Metschnikowia pulcherrima* strain to be used as a biofungicide for postharvest disease control. *Can J Microbiol* 56:128–137. <https://doi.org/10.1139/w09-117>
- Spadaro D, Ciavarella AA, Lopez-Reyes JG, Garibaldi A, Gullino ML (2010b) Effect of culture age, protectants, and initial cell concentration on viability of freeze-dried cells of *Metschnikowia pulcherrima*. *Can J Microbiol* 56:809–815. <https://doi.org/10.1139/w10-068>
- Sun C, Fu D, Lu H, Zhang J, Zheng X, Yu T (2018) Autoclaved yeast enhances the resistance against *Penicillium expansum* in postharvest pear fruit and its possible mechanisms of action. *Biol Control* 119:51–58. <https://doi.org/10.1016/j.biocontrol.2018.01.010>
- Suzuki C, Nikkuni S (1994) The primary and subunit structure of a novel type killer toxin produced by a halotolerant yeast, *Pichia farinosa*. *J Biol Chem* 269:3041–3046
- Szczęsna-Antczak M, Kamińska J, Florczak T, Turkiewicz M (2014) Cold-active yeast lipases: recent issues and future prospects. In: Buzzini P, Margesin R (eds) *Cold-adapted yeasts: biodiversity, adaptation strategies and biotechnological significance*. Springer, Berlin, pp 353–375. https://doi.org/10.1007/978-3-642-39681-6_16
- Takesako K et al (1991) Aureobasidins, new antifungal antibiotics. Taxonomy, fermentation, isolation, and properties. *J Antibiot (Tokyo)* 44:919–924. <https://doi.org/10.7164/antibiotics.44.919>
- Teichmann B, Labbe C, Lefebvre F, Bolker M, Linne U, Belanger RR (2011) Identification of a biosynthesis gene cluster for flocculosin a cellobiose lipid produced by the biocontrol agent *Pseudozyma flocculosa*. *Mol Microbiol* 79:1483–1495. <https://doi.org/10.1111/j.1365-2958.2010.07533.x>
- Tian SP, Qin GZ, Xu Y, Wang YS (2004) Application of antagonistic yeasts under field conditions and their biocontrol ability against postharvest diseases of sweet cherry. *Acta Bot Sin* 46:1324–1330
- Tilocca B, Balmás V, Hassan ZU, Jaoua S, Migheli Q (2019) A proteomic investigation of *Aspergillus carbonarius* exposed to yeast volatiles or to its major component 2-phenylethanol reveals major shifts in fungal metabolism. *Int J Food Microbiol* 306:108265. <https://doi.org/10.1016/j.ijfoodmicro.2019.108265>
- Torres R, Usall J, Teixido N, Abadias M, Viñas I (2003) Liquid formulation of the biocontrol agent *Candida sake* by modifying water activity or adding protectants. *J Appl Microbiol* 94:330–339. <https://doi.org/10.1046/j.1365-2672.2003.01843.x>
- Torres R, Teixido N, Viñas I, Mari M, Casalini L, Giraud M, Usall J (2006) Efficacy of *Candida sake* CPA-1 formulation for controlling *Penicillium expansum* decay on pome fruit from different Mediterranean regions. *J Food Prot* 69:2703–2711
- Tsai PW, Yang CY, Chang HT, Lan CY (2011) Human antimicrobial peptide LL-37 inhibits adhesion of *Candida albicans* by interacting with yeast cell-wall carbohydrates. *PLoS ONE* 6:e17755. <https://doi.org/10.1371/journal.pone.0017755>
- Turkel S, Korukluoglu M, Yavuz M (2014) Biocontrol activity of the local strain of *Metschnikowia pulcherrima* on different postharvest pathogens. *Biotechnol Res Int.* <https://doi.org/10.1155/2014/397167>
- Urquhart EJ, Punja ZK (2002) Hydrolytic enzymes and antifungal compounds produced by *Tilletiopsis* species, phyllosphere yeasts that are antagonists of powdery mildew fungi. *Can J Microbiol* 48:219–229. <https://doi.org/10.1139/w02-008>
- Usall J, Teixido N, Fons E, Viñas I (2000) Biological control of blue mould on apple by a strain of *Candida sake* under several controlled atmosphere conditions. *Int J Food Microbiol* 58:83–92. [https://doi.org/10.1016/S0168-1605\(00\)00285-3](https://doi.org/10.1016/S0168-1605(00)00285-3)
- Usall J, Torres R, Teixido N (2016) Biological control of postharvest diseases on fruit: a suitable alternative? *Curr Opin Food Sci* 11:51–55. <https://doi.org/10.1016/j.cofs.2016.09.002>
- Vadkertiova R, Molnarova J, Vranova D, Slavikova E (2012) Yeasts and yeast-like organisms associated with fruits and blossoms of different fruit trees. *Can J Microbiol* 58:1344–1352. <https://doi.org/10.1139/cjm-2012-0468>
- Vepstaitė-Monstavičė I et al (2018) *Saccharomyces paradoxus* K66 killer system evidences expanded assortment of helper and satellite viruses. *Viruses*. <https://doi.org/10.3390/v10100564>
- Verstrepen KJ, Klis FM (2006) Flocculation, adhesion and biofilm formation in yeasts. *Mol Microbiol* 60:5–15. <https://doi.org/10.1111/j.1365-2958.2006.05072.x>
- Vial L, Groleau MC, Dekimpe V, Deziel E (2007) *Burkholderia* diversity and versatility: an inventory of the extracellular products. *J Microbiol Biotechnol* 17:1407–1429
- Wachowska U, Głowacka K, Mikołajczyk W, Kucharska K (2016) Biofilm of *Aureobasidium pullulans* var. *pullulans* on winter wheat kernels and its effect on other microorganisms. *Microbiology* 85:523–530. <https://doi.org/10.1134/S0026261716050192>
- Wang X, Chi Z, Yue L, Li J (2007) Purification and characterization of killer toxin from a marine yeast *Pichia anomala* YF07b against the pathogenic yeast in crab. *Curr Microbiol* 55:396–401. <https://doi.org/10.1007/s00284-007-9010-y>
- Wang W, Chi Z, Liu G, Buzdar MA, Chi Z, Gu Q (2009a) Chemical and biological characterization of siderophore produced by the marine-derived *Aureobasidium pullulans* HN6.2 and its antibacterial activity. *Biometals* 22:965–972. <https://doi.org/10.1007/s10534-009-9248-x>

- Wang WL, Chi ZM, Chi Z, Li J, Wang XH (2009b) Siderophore production by the marine-derived *Aureobasidium pullulans* and its antimicrobial activity. *Bioresour Technol* 100:2639–2641. <https://doi.org/10.1016/j.biortech.2008.12.010>
- Wang XX, Chi Z, Peng Y, Wang XH, Ru SG, Chi ZM (2012a) Purification, characterization and gene cloning of the killer toxin produced by the marine-derived yeast *Williopsis saturnus* WC91-2. *Microbiol Res* 167:558–563. <https://doi.org/10.1016/j.micres.2011.12.001>
- Wang Y, Wei A, Li H (2012b) Using *Candida oleophila* as a biocontrol agents to prevent foodborne *Escherichia coli* O157 EHEC infections. *Springerplus* 1:82. <https://doi.org/10.1186/2193-1801-1-82>
- Weiler F, Schmitt MJ (2003) Zygocin, a secreted antifungal toxin of the yeast *Zygosaccharomyces bailii*, and its effect on sensitive fungal cells. *FEMS Yeast Res* 3:69–76. [https://doi.org/10.1016/s1567-1356\(02\)00126-5](https://doi.org/10.1016/s1567-1356(02)00126-5)
- Weiss A, Mögel G (2006) Kunz S Development of “Boni-Protect”- a yeast preparation for use in the control of post-harvest diseases of apples. In: Boos M (ed) 12th International conference on cultivation technique and phytopathological problems in organic fruit-growing. Weinsberg, Germany, pp 113–117
- Weiss A, Weisshaupt S, Krawiec P, Kunz S (2014) Use of *Aureobasidium pullulans* for resistance management in chemical control of *Botrytis cinerea* in berries. *Acta Hort* 1017:237–242. <https://doi.org/10.17660/ActaHortic.2014.1017.30>
- Wilson C, El-Ghaouth A (2002) Biological coating with a protective and curative effect for the control of postharvest decay. USA Patent US 6,423,310 B1, 23 July 2002
- Wisniewski M, Droby S (2012) Biopreservation of food and feed by postharvest biocontrol with microorganisms. In: Sundh I, Wilcks A, Goettel MS (eds) Beneficial microorganisms in agriculture, food and the environment. CABI International, Oxfordshire, pp 57–66
- Wisniewski M, Biles C, Droby S, R, Wilson C, Chalutz E (1991) Mode of action of the postharvest biocontrol yeast, *Pichia guilliermondii*. I. Characterization of attachment to *Botrytis cinerea*. *Physiol Mol Plant* 39:245–258. [https://doi.org/10.1016/0885-5765\(91\)90033-e](https://doi.org/10.1016/0885-5765(91)90033-e)
- Wisniewski M, Droby S, Chalutz E, Eilam Y (1995) Effects of Ca^{2+} and Mg^{2+} on *Botrytis cinerea* and *Penicillium expansum* *in vitro* and on the biocontrol activity of *Candida oleophila*. *Plant Pathol* 44:1016–1024. <https://doi.org/10.1111/j.1365-3059.1995.tb02660.x>
- Wisniewski M et al (2003) Characterization of a defensin in bark and fruit tissues of peach and antimicrobial activity of a recombinant defensin in the yeast, *Pichia pastoris*. *Physiol Plant* 119:563–572. <https://doi.org/10.1046/j.1399-3054.2003.00204.x>
- Wisniewski M, Wilson C, Droby S, Chalutz E, El Ghaouth A, Stevens C (2007) Postharvest biocontrol: new concepts and applications. <https://doi.org/10.1079/9781845932657.0262>
- Xu H, Nobile CJ, Dongari-Bagtzoglou A (2013) Glucanase induces filamentation of the fungal pathogen *Candida albicans*. *PLoS ONE* 8:e63736. <https://doi.org/10.1371/journal.pone.0063736>
- Yan F, Xu S, Chen Y, Zheng X (2014) Effect of rhamnolipids on *Rhotorula glutinis* biocontrol of *Alternaria alternata* infection in cherry tomato fruit. *Postharvest Biol Technol* 97:32–35. <https://doi.org/10.1016/j.postharvbio.2014.05.017>
- Yehuda H, Droby S, Wisniewski M, Goldway M (2001) A transformation system for the biocontrol yeast, *Candida oleophila*, based on hygromycin B resistance. *Curr Genet* 40:282–287. <https://doi.org/10.1007/s00294-001-0255-x>
- Yehuda H, Droby S, Bar-Shimon M, Wisniewski M, Goldway M (2003) The effect of under- and overexpressed CoEXG1-encoded exoglucanase secreted by *Candida oleophila* on the biocontrol of *Penicillium digitatum*. *Yeast* 20:771–780. <https://doi.org/10.1002/yea.1006>
- Yu T, Zheng XD (2006) Salicylic acid enhances biocontrol efficacy of the antagonist *Cryptococcus laurentii* in apple fruit. *J Plant Growth Regul* 25:166–174. <https://doi.org/10.1007/s00344-005-0077-z>
- Yurkov AM (2018) Yeasts of the soil - obscure but precious. *Yeast* 35:369–378. <https://doi.org/10.1002/yea.3310>
- Zain M, Awaad A, Razzak A, Maitland D, El-Sayed N, Sakhawy M (2009) Secondary metabolites of *Aureobasidium pullulans* isolated from Egyptian soil and their biological activity. *J Appl Sci Res* 5:1582–1591
- Zajc J, Gostincar C, Cernosa A, Gunde-Cimerman N (2019) Stress-tolerant yeasts: opportunistic pathogenicity versus biocontrol potential. *Genes (Basel)* 10:42. <https://doi.org/10.3390/genes10010042>
- Zha D, Xu L, Zhang H, Yan Y (2014) Molecular identification of lipase LipA from *Pseudomonas protegens* Pf-5 and characterization of two whole-cell biocatalysts Pf-5 and Top10lipA. *J Microbiol Biotechnol* 24:619–628. <https://doi.org/10.4014/jmb.1312.12005>
- Zhang HY, Zheng XD, Wang L, Li SS, Liu RF (2007a) Effect of yeast antagonist in combination with hot water dips on postharvest *Rhizopus* rot of strawberries. *J Food Eng* 78:281–287. <https://doi.org/10.1016/j.jfoodeng.2005.09.027>
- Zhang HY, Zheng XD, Yu T (2007b) Biological control of postharvest diseases of peach with *Cryptococcus laurentii*. *Food Control* 18:287–291. <https://doi.org/10.1016/j.foodcont.2005.10.007>
- Zhang S, Schisler DA, Boehm MJ, Slininger PJ (2007c) Utilization of chemical inducers of resistance and *Cryptococcus flavescens* OH 182.9 to reduce *Fusarium* head blight under greenhouse conditions. *Biol Control* 42:308–315. <https://doi.org/10.1016/j.biocntrol.2007.05.020>
- Zhang D, Spadaro D, Garibaldi A, Gullino ML (2011) Potential biocontrol activity of a strain of *Pichia guilliermondii* against grey mold of apples and its possible modes of action. *Biol Control* 57:193–201. <https://doi.org/10.1016/j.biocontrol.2011.02.011>
- Zhang D, Spadaro D, Valente S, Garibaldi A, Gullino ML (2012) Cloning, characterization, expression and antifungal activity of an alkaline serine protease of *Aureobasidium pullulans* PL5 involved in the biological control of postharvest pathogens. *Int J Food Microbiol* 153:453–464. <https://doi.org/10.1016/j.ijfoodmicro.2011.12.016>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.